





---

---

## Ordering properties of the smallest and largest order statistics from exponentiated location-scale models under random shocks

---

---

- Authors: MOLOD ABDOLAHI   
– Department of Statistics, Faculty of Mathematical Sciences and Computer,  
Shahid Chamran University of Ahvaz, Ahvaz, Iran  
molodabdolahi61@gmail.com
- GHOLAMALI PARHAM    
– Department of Statistics, Faculty of Mathematical Sciences and Computer,  
Shahid Chamran University of Ahvaz, Ahvaz, Iran  
parham\_g@scu.ac.ir
- RAHIM CHINIPARDAZ   
– Department of Statistics, Faculty of Mathematical Sciences and Computer,  
Shahid Chamran University of Ahvaz, Ahvaz, Iran  
chinipardaz\_r@scu.ac.ir

Received: Month 0000      Revised: Month 0000      Accepted: Month 0000

### Abstract:

- In this paper, we discuss stochastic comparisons of lifetimes of series and parallel systems when the components are exponentiated location-scale models under random shocks. The results established here are developed in two directions. First, the comparisons are carried out with respect to usual stochastic ordering by using the concept of vector majorization for series and parallel systems. Next, when the matrix of parameters changes to another matrix of parameters in the sense of multivariate chain majorization, we study the usual stochastic order of the smallest order statistics when each component receives a random shock.

### Key-Words:

- *Exponentiated location-scale family; matrix majorization; random shock; usual stochastic order; vector majorization.*

### AMS Subject Classification:

- 60E15, 60K10.

---

\*Corresponding author

---

## 1. INTRODUCTION

---

In actuarial science, it is often of interest to compare stochastically extreme claim amounts from heterogeneous portfolios. In this regard, in the present work, we compare the extreme order statistics arising from two heterogeneous portfolios in the sense of the usual stochastic ordering. It is assumed that the portfolios belong to the general exponentiated location-scale model. The general exponentiated location-scale model includes several important statistical models such as generalized exponential distribution, generalized Weibull distribution, generalized Pareto distribution and many more. The exponentiated location-scale model has three types of parameters: location, scale and shape(or skewness) parameters. Location parameter is useful in modeling the insurance related data, since an insurance company suffers a claim from policyholder after a certain period of time from the date of beginning of the policy. Also, most of the data dealing with health care costs and economy are skewed. In finance, an investigator may often have small gains, but occasionally may have a few large losses. In this case, the data will invariably be negatively skewed. If, however, we have a reverse situation, the data in this case will be positively skewed. Thus, fitting skewed models to these types of data in finance and some other fields is a very important issue. In order to fit a skewed data, we require a model having a skewness parameter. The general exponentiated location-scale model possesses this kind of flexibility. For this reason, such a general model is useful for fitting various kinds of data sets. In practical situations, the extreme claim amounts play an important role as these provide useful information for determining annual premium. In actuarial science, it is an important issue in expressing preferences between random future gains or losses. In this direction, comparisons of claim amounts in two heterogeneous portfolio of risks based on different stochastic ordering such as usual become very useful. Order statistics have received a great amount of attention from various authors. It plays an important role in several areas of probability and statistics such as reliability theory, queueing theory, and survival analysis. Let  $X_{1:n} \leq \dots \leq X_{n:n}$  denote the order statistics corresponding to the random variables  $X_1, \dots, X_n$ , where  $X_{1:n}$  and  $X_{n:n}$  corresponds to the sample minimum and sample maximum, respectively. The results of stochastic comparisons of the order statistics with independent and dependent sampling units can be seen in Dykstra et al. (1997), Zhao and Balakrishnan (2011), Li and Li (2015), Torrado (2015), Torrado and Kochar (2015), Kochar and Torrado (2015), Kundu et al. (2016), Kundu and Chowdhury (2016, 2018, 2019, 2020), Chowdhury and Kundu (2017, 2018), Hazra et al. (2017), Fang and Zhang (2013), Fang and Xu (2021), Das et al. (2019), Kundu et al. (2021), Chowdhury et al. (2021), and the references there in for a variety of parametric models. The assumption in the papers lies in the fact that each of the order statistics  $X_{1:n}, X_{2:n}, \dots, X_{n:n}$  occurs with certainty and the comparison is carried out on the minimums or the maximums of the order statistics. Now, it may so happen that the order statistics experience random shocks which may or may not result in its occurrence and it is of interest to compare two such systems stochastically with respect to vector

or matrix majorization. A random variable  $X$  is said to follow the exponentiated location-scale model if its cumulative distribution function is given by

$$(1.1) \quad F_X(x; \lambda, \theta, \alpha) = \left[ F \left( \frac{x - \lambda}{\theta} \right) \right]^\alpha, x > \lambda,$$

where  $\lambda \in R$ ,  $\alpha > 0$ ,  $\theta > 0$  and  $F$  is the baseline distribution function. Here,  $\lambda, \theta$  and  $\alpha$  are respectively known as the location, scale, and shape parameters. We write  $X \sim ELS(\lambda, \theta, \alpha)$  if  $X$  has the distribution function given by (1.1). The probability density function of the exponentiated location-scale model with (1.1) is denoted by  $f_X$ . In particular, when  $\alpha = 1$ , the model given in (1.1) reduces to the location-scale family of distributions. Further, when  $\alpha = 1$  and  $\lambda = 0$ , (1.1) reduces to the scale family and when  $\alpha = 1$  and  $\theta = 1$ , (1.1) becomes location family. The model in (1.1) coincides with the exponentiated-scale family when the location parameter  $\lambda$  is equal to 0. Thus (1.1) is a general family of distribution with great flexibility.

Let us assume series and parallel systems consist of  $n$  independent components in working conditions. Each component of the system receives a shock which may cause the component to fail. Let the random variable  $X_i$  denote lifetime of the  $i$ th component in the system which experiences a random shock at binging. Also, suppose that  $I_{p_i}$  denotes independent Bernoulli random variables, independent of  $X_i$ 's, with  $E(I_{p_i}) = p_i$ , where  $p_i$  will be called shock parameter hereafter. Then, the random shock impacts the  $i$ th component ( $I_{p_i} = 1$ ) with probability  $p_i$  or doesn't impact the  $i$ th component ( $I_{p_i} = 0$ ) with probability  $1 - p_i$ . Hence, the random variable  $Y_i = I_{p_i}X_i$  corresponds to the lifetime of  $i$ th component in a system under shock. Fang and Balakrishnan (2018) have compared two such systems with generalized Birnbaum–Saunders components. Similar comparisons are carried out in the context of the insurance where largest or smallest claim amounts in a portfolio of risks are compared stochastically. One may refer to Barmalzan et al. (2017), and Balakrishnan et al. (2018).

The survival function of  $Y_{1:n} = \min\{Y_1, \dots, Y_n\}$  is given by

$$(1.2) \quad \bar{F}_{Y_{1:n}}(x; \underline{p}, \underline{\lambda}, \underline{\theta}, \underline{\alpha}) = \prod_{i=1}^n p_i \left[ 1 - F^{\alpha_i} \left( \frac{x - \lambda_i}{\theta_i} \right) \right], x > \max\{\lambda_i, i = 1, \dots, n\},$$

and the cumulative distribution function of  $Y_{n:n} = \max\{Y_1, \dots, Y_n\}$  is given by

$$(1.3) \quad F_{Y_{n:n}}(x; \underline{p}, \underline{\lambda}, \underline{\theta}, \underline{\alpha}) = \prod_{i=1}^n \left[ 1 - p_i \left[ 1 - F^{\alpha_i} \left( \frac{x - \lambda_i}{\theta_i} \right) \right] \right], x > \max\{\lambda_i, i = 1, \dots, n\},$$

where  $\underline{x} = (x_1, \dots, x_n) \in I^n$  be any real vector and  $I^n$  denote a  $n$ -dimensional Euclidean space where  $I \subseteq R$ . Hereafter, we assume that  $Y_{1:n}^*(Y_{n:n}^*)$  denotes similarly the smallest (largest) order statistic arising from  $Y_i^* = I_{p_i^*}X_i^*$ ,  $i = 1, \dots, n$ , where  $X_1^*, \dots, X_n^*$  are independent non-negative random variables with  $X_i^* \sim ELS(\lambda_i^*, \theta_i^*, \alpha_i^*)$ ,  $i = 1, \dots, n$  and  $I_{p_1^*}, \dots, I_{p_n^*}$  are independent Bernoulli

random variables, independent of  $X_i^*$ 's, with  $E(I_{p_i^*}) = p_i^*$ ,  $i = 1, \dots, n$ . Let

$$P_n = \left\{ (\underline{x}, \underline{y}; n) : x_i > 0, y_j > 0 \text{ and } (x_i - x_j)(y_i - y_j) \leq 0, i, j = 1, \dots, n \right\},$$

$$S_n = \left\{ (\underline{x}, \underline{y}, \underline{z}; n) : x_i, y_j, z_k > 0 \text{ and } x_i \leq (\geq) x_j, y_i \geq (\leq) y_j, z_i \geq (\leq) z_j \right\},$$

$$N_n = \left\{ (\underline{x}, \underline{y}, \underline{z}; n) : z_i \geq 1, x_i > 0, y_i > 0, x_i \leq (\geq) x_j, y_i \leq (\geq) y_j, z_i \leq (\geq) z_j \right\},$$

$$N_n^* = \left\{ (\underline{x}, \underline{y}, \underline{z}; n) : z_i > 0, x_i > 0, y_i > 0, x_i \leq (\geq) x_j, y_i \leq (\geq) y_j, z_i \leq (\geq) z_j \right\},$$

$$U_n = \left\{ (\underline{w}, \underline{x}, \underline{y}, \underline{z}; n) : w_i, x_j, y_k, z_l > 0, w_i \leq (\geq) w_j, x_i \geq (\leq) x_j, y_i \geq (\leq) y_j, \right. \\ \left. z_i \geq (\leq) z_j \right\}.$$

The rest of this paper is organized as follows. In Section 2, we introduce some definitions and fundamental lemmas. In Section 3, we establish some ordering properties for the smallest and largest order statistics of the ELS model with associated random shocks. In Section 4, some special cases of our main results are added. Section 5 provides applications of the established results. Finally, in Section 6, we include some concluding.

---

## 2. PRELIMINARIES

---

In this section we provide some preliminary definitions and lemmas which will be useful in the sequel. To compare lifetimes of series and parallel systems, stochastic orders have been extensively used in the literature. Below, we present a few of them. Throughout the paper, we use the notations  $R = (-\infty, +\infty)$ ,  $R_+ = (0, +\infty)$  and  $a \stackrel{\text{sign}}{=} b$  means that  $a$  and  $b$  have the same sign. Let  $X$  be non-negative random variable with distribution function  $F$ , and density function  $f$ . The survival function, hazard rate, and reversed hazard rate are  $\bar{F} = 1 - F$ ,  $r_X = \frac{f}{F}$ , and  $\tilde{r}_X = \frac{f}{\bar{F}}$ , respectively.

**Definition 2.1.** Let  $X$  and  $Y$  be two absolutely continuous random variables with respective supports  $(l_X, u_X)$  and  $(l_Y, u_Y)$ , where  $u_X$  and  $u_Y$  may be positive infinity, and  $l_X$  and  $l_Y$  may be negative infinity. Then,  $X$  is said to be smaller than  $Y$  in usual stochastic (st) order, denoted as  $X \leq_{st} Y$ , if  $\bar{F}_X(t) \leq \bar{F}_Y(t)$  for all  $t \in (-\infty, +\infty)$ .

Let  $\underline{x} = (x_1, \dots, x_n) \in I^n$  and  $\underline{y} = (y_1, \dots, y_n) \in I^n$  be any two real vectors with  $x_{(1)}, \dots, x_{(n)}$  being the increasing arrangement of the components of the vector  $\underline{x}$ . The following definitions may be found in Marshall et al. (2011).

**Definition 2.2.** (i) The vector  $\underline{x}$  is said to majorize the vector  $\underline{y}$  (written as  $\underline{x} \stackrel{m}{\geq} \underline{y}$ ) if

$$\sum_{i=1}^j x_{(i)} \leq \sum_{i=1}^j y_{(i)}, j = 1, \dots, n-1, \text{ and } \sum_{i=1}^n x_{(i)} = \sum_{i=1}^n y_{(i)};$$

(ii) The vector  $\underline{x}$  is said to weakly supermajorize the vector  $\underline{y}$  (written as  $\underline{x} \stackrel{w}{\geq} \underline{y}$ ) if

$$\sum_{i=1}^j x_{(i)} \leq \sum_{i=1}^j y_{(i)} \text{ for } j = 1, \dots, n;$$

(iii) The vector  $\underline{x}$  is said to weakly submajorize the vector  $\underline{y}$  (written as  $\underline{x} \stackrel{w}{\geq} \underline{y}$ ) if

$$\sum_{i=j}^n x_{(i)} \geq \sum_{i=j}^n y_{(i)} \text{ for } j = 1, \dots, n;$$

(iv) The vector  $\underline{x}$  is said to be p-larger than the vector  $\underline{y}$  (written as  $\underline{x} \stackrel{p}{\geq} \underline{y}$ ) if

$$\prod_{i=1}^j x_{(i)} \leq \prod_{i=1}^j y_{(i)} \text{ for } j = 1, \dots, n;$$

(v) The vector  $\underline{x}$  is said to reciprocally majorize the vector  $\underline{y}$  (written as  $\underline{x} \stackrel{rm}{\geq} \underline{y}$ ) if

$$\sum_{i=1}^j \frac{1}{x_{(i)}} \geq \sum_{i=1}^j \frac{1}{y_{(i)}} \text{ for } j = 1, \dots, n.$$

It is not difficult to show that  $\underline{x} \stackrel{m}{\geq} \underline{y} \Rightarrow \underline{x} \stackrel{w}{\geq} \underline{y} \Rightarrow \underline{x} \stackrel{p}{\geq} \underline{y} \Rightarrow \underline{x} \stackrel{rm}{\geq} \underline{y}$ .

**Definition 2.3.** A function  $\psi : I^n \rightarrow R$  is said to be schur-convex (schur-concave) on  $I^n$  if

$$\underline{x} \stackrel{m}{\geq} \underline{y} \text{ implies } \psi(\underline{x}) \geq (\leq) \psi(\underline{y}) \text{ for all } \underline{x}, \underline{y} \in I^n.$$

The following definitions related to matrix majorization may be found in Marshall et al. (2011).

**Definition 2.4.** (i) A square matrix  $\Pi_n$ , of order  $n$ , is said to be a permutation matrix if each row and column has a single entry 1, and all other entries as zero;

(ii) A square matrix  $P = (p_{ij})$ , of order  $n$ , is said to be doubly stochastic if  $p_{ij} \geq 0$ , for all  $i, j = 1, \dots, n$ ,  $\sum_{i=1}^n p_{ij} = 1$ ,  $j = 1, \dots, n$  and  $\sum_{j=1}^n p_{ij} = 1$ ,  $i = 1, \dots, n$ ;

(iii) A square matrix  $T_w$ , is said to be T-transform matrix if it has form  $T_w = wI + (1-w)\Pi$ ; where  $0 \leq w \leq 1$ ,  $I$  is the identity matrix and  $\Pi$  is the permutation matrix. Let  $T_{w_1} = w_1I + (1-w_1)\Pi_1$  and  $T_{w_2} = w_2I + (1-w_2)\Pi_2$  be two transform matrices, where  $0 \leq w_1, w_2 \leq 1$  and  $\Pi_1$  and  $\Pi_2$  are two permutation matrices that interchange two coordinates. Then, we say  $T_{w_1}$  and  $T_{w_2}$  have the same structure if  $\Pi_1 = \Pi_2$ , where  $\Pi_1$  and  $\Pi_2$  are permutation matrices with the same dimension, otherwise they are different structures.

**Definition 2.5.** Consider the  $m \times n$  matrices  $A = \{a_{ij}\}$  and  $B = \{b_{ij}\}$  with rows  $a_1, \dots, a_m$  and  $b_1, \dots, b_n$ , respectively.

(i)  $A$  is said to be larger than  $B$  in chain majorization, denoted by  $A \gg B$ , if there exists a finite set of  $n \times n$  T-transform matrices  $T_1, \dots, T_k$  such that  $B = AT_1 \cdots T_k$ ;

(ii)  $A$  is said to majorize  $B$ , denoted by  $A > B$ , if  $A = BP$ , where the  $n \times n$  matrix  $P$  is doubly stochastic. Since a product of T-transforms is doubly stochastic, it follows that  $A \gg B \Rightarrow A > B$ ;

(iii)  $A$  is said to be larger than the matrix  $B$  in row majorization, denoted by  $A \overset{row}{>} B$ , if  $a_i \geq b_i$  for  $i = 1, \dots, m$ . It is clear that  $A > B \Rightarrow A \overset{row}{>} B$ ;

(iv)  $A$  is said to be larger than the matrix  $B$  in row weakly majorization, denoted by  $A \overset{w}{>} B$ , if  $a_i \overset{w}{\geq} b_i$  for  $i = 1, \dots, m$ . It is clear that  $A \overset{row}{>} B \Rightarrow A \overset{w}{>} B$ . Thus it can be written that  $A \gg B \Rightarrow A > B \Rightarrow A \overset{row}{>} B \Rightarrow A \overset{w}{>} B$ .

Also, we introduce the following notations.

(i)  $D_+ = \{(x_1, \dots, x_n) : x_1 \geq \dots \geq x_n > 0\}$ ;

(ii)  $\varepsilon_+ = \{(x_1, \dots, x_n) : 0 < x_1 \leq \dots \leq x_n\}$ ;

(iii)  $(h(\underline{p}), \underline{\lambda}) = \begin{pmatrix} h(p_1) & h(p_2) & \cdots & h(p_n) \\ \lambda_1 & \lambda_2 & \cdots & \lambda_n \end{pmatrix}$ .

The following two Lemmas are used to prove the two Theorems 2.17 and 2.18.

**Lemma 2.1.** A differentiable function  $\Phi : R_+^8 \rightarrow R_+$  satisfies

$$(2.1) \quad \Phi(A) \geq (\leq) \Phi(B) \text{ for all } A, B \text{ such that } A \in U_2, A \gg B$$

if and only if

(i)  $\Phi(A) = \Phi(A\Pi)$  for all permutation matrix  $\Pi$  and for all  $A \in U_2$ , and

(ii)  $\sum_{i=1}^4 (a_{ik} - a_{ij}) [\Phi_{ik}(A) - \Phi_{ij}(A)] \geq (\leq) 0$  for all  $j, k = 1, 2$  and for all  $A \in U_2$ ,

where  $\Phi_{ij}(A) = \frac{\partial \Phi(A)}{\partial a_{ij}}$ .

**Lemma 2.2.** *Let the function  $\gamma : R_+^4 \rightarrow R_+$  be differentiable and the function  $\Phi_n : R_+^{4n} \rightarrow R_+$  be defined as*

$$\Phi_n(A) = \prod_{i=1}^n \gamma(a_{1i}, a_{2i}, a_{3i}, a_{4i}).$$

Assume that  $\Phi_2$  satisfies (2.1). Then for  $A \in U_n$  and  $B = AT_w$ , we have  $\Phi_n(A) \geq (\leq) \Phi_n(B)$ , where  $T_w$  is the  $T$ -transform matrix.

**Proof:** The proofs of Lemmas 2.1-2.2 are similar to those of Theorems 2 and 3 Balakrishnan et al. (2015), and Marshall and Olkin (1997).  $\square$

---

### 3. MAIN RESULTS

---

In this section we establish some ordering properties for the smallest and largest order statistics of the ELS model with associated random shocks. We now consider the following assumption.

**Assumption 3.1** Suppose  $X_1, \dots, X_n$  are independent non-negative random variables with  $X_i \sim ELS(\lambda_i, \theta_i, \alpha_i)$ , and  $I_{p_1}, \dots, I_{p_n}$  are independent Bernoulli random variables, independent of  $X_i$ 's, with  $E(I_{p_i}) = p_i, i = 1, \dots, n$ . Further, suppose  $X_1^*, \dots, X_n^*$  are independent non-negative random variables with  $X_i^* \sim ELS(\lambda_i^*, \theta_i^*, \alpha_i^*)$ , and  $I_{p_1}^*, \dots, I_{p_n}^*$  are independent Bernoulli random variables, independent of  $X_i^*$ 's, with  $E(I_{p_i}^*) = p_i^*, i = 1, \dots, n$ .

Theorem 3.1 shows that usual stochastic ordering holds between two parallel systems of heterogeneous components under random shocks for fixed  $\theta$  and  $\alpha$ .

**Theorem 3.1.** *Let Assumption 3.1 hold and  $h : [0, 1] \rightarrow R$  be a differentiable, increasing and strictly convex function. Also,  $\lambda_i = \lambda_i^*, \theta_i = \theta_i^* = \theta$  and  $\alpha_i = \alpha_i^* = \alpha$ , where  $i = 1, \dots, n$ . Then,  $h(\underline{p}) \geq_w h(\underline{p}^*)$  implies  $Y_{n:n} \geq_{st} Y_{n:n}^*$ , provided  $\underline{\lambda} \in D_+, h(\underline{p}) \in D_+$ , and  $h(\underline{p}) = (h(p_1), \dots, h(p_n))$ .*

**Proof:** The cumulative distribution function of  $Y_{n:n}$  is given by

$$F_{Y_{n:n}}(x) = \prod_{i=1}^n \left[ 1 - h^{-1}(u_i) \left[ 1 - F^\alpha \left( \frac{x - \lambda_i}{\theta} \right) \right] \right],$$

where  $h(p_i) = u_i$ . Let us define  $\psi_1(\underline{y}) = F_{Y_{n:n}}(x)$ . Differentiating  $\psi_1(\underline{y})$ , partially, with respect to  $u_i$ , we get

$$(3.1) \quad \frac{\partial \psi_1(\underline{y})}{\partial u_i} = - \frac{\frac{dh^{-1}(u_i)}{du_i} \left( 1 - F^\alpha \left( \frac{x - \lambda_i}{\theta} \right) \right)}{1 - h^{-1}(u_i) \left( 1 - F^\alpha \left( \frac{x - \lambda_i}{\theta} \right) \right)} \psi_1(\underline{y}) \leq 0,$$

so,  $\psi_1(\underline{y})$  is decreasing in each  $u_i$ . Again, it can be shown that

$$(3.2) \quad \frac{\partial \psi_1(\underline{y})}{\partial u_i} - \frac{\partial \psi_1(\underline{y})}{\partial u_j} \stackrel{\text{sign}}{\leq} \frac{\frac{dh^{-1}(u_j)}{du_j} \left(1 - F^\alpha \left(\frac{x-\lambda_j}{\theta}\right)\right)}{1 - h^{-1}(u_j) \left(1 - F^\alpha \left(\frac{x-\lambda_j}{\theta}\right)\right)} - \frac{\frac{dh^{-1}(u_i)}{du_i} \left(1 - F^\alpha \left(\frac{x-\lambda_i}{\theta}\right)\right)}{1 - h^{-1}(u_i) \left(1 - F^\alpha \left(\frac{x-\lambda_i}{\theta}\right)\right)}$$

Now,

$$\frac{\partial}{\partial u} \left( \frac{1}{1 - h^{-1}(u) \left(1 - F^\alpha \left(\frac{x-\lambda}{\theta}\right)\right)} \right) = \frac{dh^{-1}(u)}{du} \left(1 - F^\alpha \left(\frac{x-\lambda}{\theta}\right)\right) \geq 0,$$

implying that  $\frac{1}{1 - h^{-1}(u) \left(1 - F^\alpha \left(\frac{x-\lambda}{\theta}\right)\right)}$  is increasing in  $u$ . Thus, as  $\lambda \in D_+$ ,  $h(p) \in$

$D_+$ , for  $i \leq j$  taking  $\lambda_i \geq \lambda_j$  and  $u_i \geq u_j$  and noticing that  $h^{-1}(u)$  is increasing in  $u$ , it can be written that

$$\frac{1 - F^\alpha \left(\frac{x-\lambda_j}{\theta}\right)}{1 - h^{-1}(u_j) \left(1 - F^\alpha \left(\frac{x-\lambda_j}{\theta}\right)\right)} \leq \frac{1 - F^\alpha \left(\frac{x-\lambda_i}{\theta}\right)}{1 - h^{-1}(u_i) \left(1 - F^\alpha \left(\frac{x-\lambda_i}{\theta}\right)\right)},$$

Again, if  $h(u)$  is convex in  $u$ , then  $u_i \geq u_j$  gives  $\frac{dh^{-1}(u_i)}{du_i} \geq \frac{dh^{-1}(u_j)}{du_j}$  which yields

$$\frac{\frac{dh^{-1}(u_j)}{du_j} \left(1 - F^\alpha \left(\frac{x-\lambda_j}{\theta}\right)\right)}{1 - h^{-1}(u_j) \left(1 - F^\alpha \left(\frac{x-\lambda_j}{\theta}\right)\right)} \leq \frac{\frac{dh^{-1}(u_i)}{du_i} \left(1 - F^\alpha \left(\frac{x-\lambda_i}{\theta}\right)\right)}{1 - h^{-1}(u_i) \left(1 - F^\alpha \left(\frac{x-\lambda_i}{\theta}\right)\right)}.$$

Substituting the above result in (3.2), we get  $\frac{\partial \psi_1(\underline{y})}{\partial u_i} - \frac{\partial \psi_1(\underline{y})}{\partial u_j} \leq 0$ . Thus, by Lemma 3.1 of Kundu et al. (2016),  $\psi_1(\underline{y})$  is Schur concave in  $\underline{y}$ . Thus the result is proved by Theorem A.8 of Marshall et al. (2011).  $\square$

Theorem 3.2 shows that the majorized shape parameter vector leads to smaller systems lifetime in the sense of the usual stochastic ordering when the location and scale parameter vectors are constant and shock parameter vectors are heterogeneous.

**Theorem 3.2.** *Let Assumption 3.1 hold and  $h : [0, 1] \rightarrow R$  be a differentiable and decreasing function. Also,  $\lambda_i = \lambda_i^* = \lambda$ ,  $\theta_i = \theta_i^* = \theta$ , and  $\alpha_i^* = \beta_i$ , where  $i = 1, \dots, n$ . Then,  $\underline{\alpha} \stackrel{w}{\geq} \underline{\beta}$  implies  $Y_{n:n} \leq_{st} Y_{n:n}^*$ , provided  $\underline{\alpha}, \underline{\beta}, h(p) \in \varepsilon_+$ .*

**Proof:** The cumulative distribution function of  $Y_{n:n}$  is given by

$$F_{Y_{n:n}}(x) = \prod_{i=1}^n \left[ 1 - h^{-1}(u_i) \left[ 1 - F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right) \right] \right],$$



where  $h(p_i) = u_i$ . Differentiating  $F_{Y_{n:n}}(x)$ , partially, with respect to  $\alpha_i$ , we get

$$\frac{\partial F_{Y_{n:n}}(x)}{\partial \alpha_i} = \frac{h^{-1}(u_i)F^{\alpha_i}\left(\frac{x-\lambda}{\theta}\right) \ln[F\left(\frac{x-\lambda}{\theta}\right)]}{1 - h^{-1}(u_i)\left(1 - F^{\alpha_i}\left(\frac{x-\lambda}{\theta}\right)\right)} F_{Y_{n:n}}(x) \leq 0,$$

so,  $F_{Y_{n:n}}(x)$  is decreasing in each  $\alpha_i$ . Again, it can be shown that

$$(3.3) \quad \frac{\partial F_{Y_{n:n}}(x)}{\partial \alpha_i} - \frac{\partial F_{Y_{n:n}}(x)}{\partial \alpha_j} \stackrel{\text{sign}}{=} \frac{h^{-1}(u_i)F^{\alpha_i}\left(\frac{x-\lambda}{\theta}\right) \ln[F\left(\frac{x-\lambda}{\theta}\right)]}{1 - h^{-1}(u_i)\left(1 - F^{\alpha_i}\left(\frac{x-\lambda}{\theta}\right)\right)} - \frac{h^{-1}(u_j)F^{\alpha_j}\left(\frac{x-\lambda}{\theta}\right) \ln[F\left(\frac{x-\lambda}{\theta}\right)]}{1 - h^{-1}(u_j)\left(1 - F^{\alpha_j}\left(\frac{x-\lambda}{\theta}\right)\right)}.$$

Now,

$$\frac{\partial}{\partial \alpha} \left( \frac{F^\alpha\left(\frac{x-\lambda}{\theta}\right)}{1 - h^{-1}(u)(1 - F^\alpha\left(\frac{x-\lambda}{\theta}\right))} \right) \stackrel{\text{sign}}{=} (1 - h^{-1}(u))F^\alpha\left(\frac{x-\lambda}{\theta}\right) \ln[F\left(\frac{x-\lambda}{\theta}\right)] \leq 0,$$

implying that  $\frac{F^\alpha\left(\frac{x-\lambda}{\theta}\right)}{1 - h^{-1}(u)\left(1 - F^\alpha\left(\frac{x-\lambda}{\theta}\right)\right)}$  is decreasing in  $\alpha$ . Again, as  $h(u)$  is decreasing in  $u$ , then

$$\frac{\partial}{\partial u} \left( \frac{h^{-1}(u)}{1 - h^{-1}(u)(1 - F^\alpha\left(\frac{x-\lambda}{\theta}\right))} \right) \stackrel{\text{sign}}{=} \frac{\partial h^{-1}(u)}{\partial u} \leq 0,$$

implying that  $\frac{h^{-1}(u)}{1 - h^{-1}(u)(1 - F^\alpha\left(\frac{x-\lambda}{\theta}\right))}$  is decreasing in  $u$ . Thus, as  $\alpha \in \varepsilon_+$ ,  $h(p) \in \varepsilon_+$ , for  $i \leq j$  taking  $\alpha_i \leq \alpha_j$  and  $u_i \leq u_j$  and noticing that  $h^{-1}(u)$  is decreasing in  $u$ , it can be written that

$$\begin{aligned} \frac{h^{-1}(u_j)F^{\alpha_j}\left(\frac{x-\lambda}{\theta}\right)}{1 - h^{-1}(u_j)\left(1 - F^{\alpha_j}\left(\frac{x-\lambda}{\theta}\right)\right)} &\leq \frac{h^{-1}(u_j)F^{\alpha_i}\left(\frac{x-\lambda}{\theta}\right)}{1 - h^{-1}(u_j)\left(1 - F^{\alpha_i}\left(\frac{x-\lambda}{\theta}\right)\right)} \\ &\leq \frac{h^{-1}(u_i)F^{\alpha_i}\left(\frac{x-\lambda}{\theta}\right)}{1 - h^{-1}(u_i)\left(1 - F^{\alpha_i}\left(\frac{x-\lambda}{\theta}\right)\right)}, \end{aligned}$$

which implies

$$\begin{aligned} \frac{h^{-1}(u_j)F^{\alpha_j}\left(\frac{x-\lambda}{\theta}\right) \ln[F\left(\frac{x-\lambda}{\theta}\right)]}{1 - h^{-1}(u_j)\left(1 - F^{\alpha_j}\left(\frac{x-\lambda}{\theta}\right)\right)} &\geq \frac{h^{-1}(u_j)F^{\alpha_i}\left(\frac{x-\lambda}{\theta}\right) \ln[F\left(\frac{x-\lambda}{\theta}\right)]}{1 - h^{-1}(u_j)\left(1 - F^{\alpha_i}\left(\frac{x-\lambda}{\theta}\right)\right)} \\ &\geq \frac{h^{-1}(u_i)F^{\alpha_i}\left(\frac{x-\lambda}{\theta}\right) \ln[F\left(\frac{x-\lambda}{\theta}\right)]}{1 - h^{-1}(u_i)\left(1 - F^{\alpha_i}\left(\frac{x-\lambda}{\theta}\right)\right)}. \end{aligned}$$

Hence, substituting the above results in (3.3), we get  $\frac{\partial F_{Y_{n:n}}(x)}{\partial \alpha_i} - \frac{\partial F_{Y_{n:n}}(x)}{\partial \alpha_j} \leq 0$ . Thus, by Lemma 3.3 of Kundu et al. (2016), it can be proved that  $F_{Y_{n:n}}(x)$  is Schur-convex in  $\alpha$ . Thus the result is proved by Theorem A.8 of Marshall et al. (2011).  $\square$

**Theorem 3.3.** *Let Assumption 3.1 hold and  $h : [0, 1] \rightarrow R$  be a differentiable and decreasing function. Also,  $\lambda_i = \lambda_i^* = \lambda$ ,  $\theta_i = \theta_i^* = \theta$ , and  $\alpha_i^* = \beta_i$ , where  $i = 1, \dots, n$ . Then,  $\underline{\alpha}^{-1} \geq_w \underline{\beta}^{-1}$  implies  $Y_{n:n} \leq_{st} Y_{n:n}^*$ , provided,  $\underline{\alpha}, \underline{\beta}, h(\underline{p}) \in \varepsilon_+$ .*

**Proof:** The cumulative distribution function of  $Y_{n:n}$  can be expressed as the function of  $a_i$ , where  $a_i = \frac{1}{\alpha_i}$ ,  $i = 1, \dots, n$ . We denote it by  $\psi_2(\underline{a})$ , where  $\underline{a} = (a_1, \dots, a_n)$ , and

$$\psi_2(\underline{a}) = \prod_{i=1}^n \left[ 1 - h^{-1}(u_i) \left[ 1 - F^{\frac{1}{a_i}} \left( \frac{x - \lambda}{\theta} \right) \right] \right],$$

where  $h(p_i) = u_i$ . It can be shown that the partial derivative of  $\psi_2(\underline{a})$ , with respect to  $a_i$  increasing in  $i = 1, \dots, n$ . Thus, by Lemma 3.1 of Kundu et al. (2016), it can be proved that  $\psi_2(\underline{a})$  is Schur-convex in  $\underline{a}$ . Thus the result is proved by Theorem A.8 of Marshall et al. (2011).  $\square$

Theorem 3.4 shows that  $Y_{n:n}$  is smaller than  $Y_{n:n}^*$  with respect to the usual stochastic ordering when a vector of scale parameters is  $p$ -larger than that of another vector of the scale parameters with some additional conditions when the location and shape parameter vectors are constant and shock parameter vectors are heterogeneous. Similar result also hold under reciprocally majorized based conditions among the scale parameters.

**Theorem 3.4.** *Let Assumption 3.1 hold and  $h : [0, 1] \rightarrow R$  be a differentiable and decreasing function. Also,  $\lambda_i = \lambda_i^* = \lambda$ ,  $\theta_i^* = \delta_i$ , and  $\alpha_i = \alpha_i^* = \alpha$ , where  $i = 1, \dots, n$ . Then,*

- (i)  $\underline{\theta} \stackrel{p}{\geq} \underline{\delta}$  implies  $Y_{n:n} \leq_{st} Y_{n:n}^*$ , provided  $\underline{\theta}, \underline{\delta}, h(\underline{p}) \in D_{+,}$ , and  $u\tilde{r}(u)$  is increasing in  $u$ ;
- (ii)  $\underline{\theta} \stackrel{rm}{\geq} \underline{\delta}$  implies  $Y_{n:n} \leq_{st} Y_{n:n}^*$ , provided  $\underline{\theta}, \underline{\delta}, h(\underline{p}) \in \varepsilon_+$ , and  $\tilde{r}(u)$  is increasing in  $u$ .

**Proof(i) :** The cumulative distribution function of  $Y_{n:n}$  can be expressed as the function of  $a_i$ , where  $a_i = \ln \theta_i$ ,  $i = 1, \dots, n$ . We denote it by  $\psi_3(\underline{a})$ , where  $\underline{a} = (a_1, \dots, a_n)$ , and

$$\psi_3(\underline{a}) = \prod_{i=1}^n \left[ 1 - h^{-1}(u_i) [1 - F^\alpha(e^{-a_i}(x - \lambda))] \right],$$

where  $h(p_i) = u_i$ . Differentiating  $\psi_3(\underline{a})$ , partially, with respect to  $a_i$ , we get

$$\frac{\partial \psi_3(\underline{a})}{\partial a_i} = - \frac{\alpha h^{-1}(u_i) e^{-a_i} (x - \lambda) \tilde{r}(e^{-a_i}(x - \lambda)) F^\alpha(e^{-a_i}(x - \lambda))}{1 - h^{-1}(u_i) (1 - F^\alpha(e^{-a_i}(x - \lambda)))} \psi_3(\underline{a}) \leq 0,$$

so,  $\psi_3(\underline{a})$  is decreasing in each  $a_j$ . Again, it can be shown that

$$(3.4) \quad \frac{\partial \psi_3(\underline{a})}{\partial a_i} - \frac{\partial \psi_3(\underline{a})}{\partial a_j} \stackrel{\text{sign}}{=} \frac{\alpha h^{-1}(u_j) e^{-a_j} (x - \lambda) \tilde{r}(e^{-a_j} (x - \lambda)) F^\alpha(e^{-a_j} (x - \lambda))}{1 - h^{-1}(u_j) (1 - F^\alpha(e^{-a_j} (x - \lambda)))} - \frac{\alpha h^{-1}(u_i) e^{-a_i} (x - \lambda) \tilde{r}(e^{-a_i} (x - \lambda)) F^\alpha(e^{-a_i} (x - \lambda))}{1 - h^{-1}(u_i) (1 - F^\alpha(e^{-a_i} (x - \lambda)))}.$$

Now,

$$\frac{\partial}{\partial a} \left( \frac{F^\alpha(e^{-a} (x - \lambda))}{1 - h^{-1}(u) (1 - F^\alpha(e^{-a} (x - \lambda)))} \right) \stackrel{\text{sign}}{=} -\alpha e^{-a} (x - \lambda) (1 - h^{-1}(u)) \times \tilde{r}(e^{-a} (x - \lambda)) \times F^\alpha(e^{-a} (x - \lambda)) \leq 0,$$

implying that  $\frac{F^\alpha(e^{-a} (x - \lambda))}{1 - h^{-1}(u) (1 - F^\alpha(e^{-a} (x - \lambda)))}$  is decreasing in  $a$ . Again, as  $h(u)$  is decreasing in  $u$ , then Fang, L. and Balakrishnan, N. (2018). Ordering properties of the small25 est order statistics from generalized Birnbaum?Saunders models with associated 26 random shocks, *Metrika*, 81, 1, 19-35.

$$\frac{\partial}{\partial u} \left( \frac{h^{-1}(u)}{1 - h^{-1}(u) (1 - F^\alpha(e^{-a} (x - \lambda)))} \right) \stackrel{\text{sign}}{=} \frac{\partial h^{-1}(u)}{\partial u} \leq 0,$$

implying that  $\frac{h^{-1}(u)}{1 - h^{-1}(u) (1 - F^\alpha(e^{-a} (x - \lambda)))}$  is decreasing in  $u$ . Thus, as  $\underline{a}, h(p) \in D_+$ , for  $i \leq j$  taking  $a_i \geq a_j$  and  $u_i \geq u_j$  and noticing that  $h^{-1}(u)$  is decreasing in  $u$ , it can be written that

$$\begin{aligned} \frac{h^{-1}(u_j) F^\alpha(e^{-a_j} (x - \lambda))}{1 - h^{-1}(u_j) (1 - F^\alpha(e^{-a_j} (x - \lambda)))} &\geq \frac{h^{-1}(u_j) F^\alpha(e^{-a_i} (x - \lambda))}{1 - h^{-1}(u_j) (1 - F^\alpha(e^{-a_i} (x - \lambda)))} \\ &\geq \frac{h^{-1}(u_i) F^\alpha(e^{-a_i} (x - \lambda))}{1 - h^{-1}(u_i) (1 - F^\alpha(e^{-a_i} (x - \lambda)))}, \end{aligned}$$

As  $u \tilde{r}(u)$  is increasing in  $u$ , then

$$\begin{aligned} \frac{\alpha h^{-1}(u_j) e^{-a_j} (x - \lambda) \tilde{r}(e^{-a_j} (x - \lambda)) F^\alpha(e^{-a_j} (x - \lambda))}{1 - h^{-1}(u_j) (1 - F^\alpha(e^{-a_j} (x - \lambda)))} &\geq \\ \frac{\alpha h^{-1}(u_i) e^{-a_i} (x - \lambda) \tilde{r}(e^{-a_i} (x - \lambda)) F^\alpha(e^{-a_i} (x - \lambda))}{1 - h^{-1}(u_i) (1 - F^\alpha(e^{-a_i} (x - \lambda)))}. \end{aligned}$$

Hence, from (3.4), we get  $\frac{\partial \psi_3(\underline{a})}{\partial a_i} - \frac{\partial \psi_3(\underline{a})}{\partial a_j} \geq 0$ . Thus, by Lemma 3.1 of Kundu et al. (2016), it can be proved that  $\psi_3(\underline{a})$  is Schur-convex in  $\underline{a}$ . Thus the result is proved by Lemma 3.1 of Khaledi et al. (2002).

**Proof (ii)** : The cumulative distribution function of  $Y_{n:n}$  can be expressed as the function of  $b_i = \frac{1}{\theta_i}$ ,  $i = 1, \dots, n$ . We denote it by  $\psi_4(\underline{b})$ , where  $\underline{b} = (b_1, \dots, b_n)$ .

$$\psi_4(\underline{b}) = \prod_{i=1}^n \left[ 1 - h^{-1}(u_i) [1 - F^\alpha(b_i(x - \lambda))] \right],$$

where  $h(p_i) = u_i$ . Differentiating  $\psi_4(\underline{b})$ , partially, with respect to  $b_i$ , we see that  $\psi_4(\underline{b})$  is increasing in each  $i = 1, \dots, n$ . Thus, by Lemma 3.1 of Kundu et al. (2016), it can be proved that  $\psi_4(\underline{b})$  is Schur-convex in  $\underline{b}$ . Thus the result is proved by Lemma 4.1 of Hazra et al. (2017).

Theorem 3.5 shows that the majorized shape parameter vector leads to smaller systems lifetime in the sense of the usual stochastic ordering when the location and scale and shock parameter vectors are heterogeneous.

**Theorem 3.5.** *Let Assumption 3.1 hold and  $h : [0, 1] \rightarrow R$  be a differentiable and decreasing function. Also,  $\lambda_i = \lambda_i^*$ ,  $\theta_i = \theta_i^*$ , and  $\alpha_i^* = \beta_i$ , where  $i = 1, \dots, n$ . Then,  $\alpha \underset{\sim}{\succ} \beta$  implies  $Y_{n:n} \leq_{st} Y_{n:n}^*$ , provided  $\alpha, \beta, h(\underline{p}), \underline{\lambda}, \underline{\theta}, \in \varepsilon_+$ .*

**Proof:** The cumulative distribution function of  $Y_{n:n}$  is given by

$$F_{Y_{n:n}}(x) = \prod_{i=1}^n \left[ 1 - h^{-1}(u_i) \left[ 1 - F^{\alpha_i} \left( \frac{x - \lambda_i}{\theta_i} \right) \right] \right],$$

where  $h(p_i) = u_i$ . Differentiating  $F_{Y_{n:n}}(x)$ , partially, with respect to  $\alpha_i$ , we get

$$\frac{\partial F_{Y_{n:n}}(x)}{\partial \alpha_i} = \frac{h^{-1}(u_i) F^{\alpha_i} \left( \frac{x - \lambda_i}{\theta_i} \right) \ln \left[ F \left( \frac{x - \lambda_i}{\theta_i} \right) \right]}{1 - h^{-1}(u_i) \left( 1 - F^{\alpha_i} \left( \frac{x - \lambda_i}{\theta_i} \right) \right)} F_{Y_{n:n}}(x) \leq 0,$$

so,  $F_{Y_{n:n}}(x)$  is decreasing in each  $\alpha_i$ . Again, it can be shown that

$$\begin{aligned} \frac{\partial F_{Y_{n:n}}(x)}{\partial \alpha_i} - \frac{\partial F_{Y_{n:n}}(x)}{\partial \alpha_j} &\stackrel{\text{sign}}{=} \frac{h^{-1}(u_i) F^{\alpha_i} \left( \frac{x - \lambda_i}{\theta_i} \right) \ln \left[ F \left( \frac{x - \lambda_i}{\theta_i} \right) \right]}{1 - h^{-1}(u_i) \left( 1 - F^{\alpha_i} \left( \frac{x - \lambda_i}{\theta_i} \right) \right)} \\ &\quad - \frac{h^{-1}(u_j) F^{\alpha_j} \left( \frac{x - \lambda_j}{\theta_j} \right) \ln \left[ F \left( \frac{x - \lambda_j}{\theta_j} \right) \right]}{1 - h^{-1}(u_j) \left( 1 - F^{\alpha_j} \left( \frac{x - \lambda_j}{\theta_j} \right) \right)}. \end{aligned}$$

Now,

$$\frac{\partial}{\partial \alpha} \left( \frac{F^\alpha \left( \frac{x - \lambda}{\theta} \right)}{1 - h^{-1}(u) \left( 1 - F^\alpha \left( \frac{x - \lambda}{\theta} \right) \right)} \right) \stackrel{\text{sign}}{=} (1 - h^{-1}(u)) F^\alpha \left( \frac{x - \lambda}{\theta} \right) \ln \left[ F \left( \frac{x - \lambda}{\theta} \right) \right] \leq 0,$$

implying that  $\frac{F^\alpha \left( \frac{x - \lambda}{\theta} \right)}{1 - h^{-1}(u) \left( 1 - F^\alpha \left( \frac{x - \lambda}{\theta} \right) \right)}$  is decreasing in  $\alpha$ . Again, as  $h(u)$  is decreasing in  $u$ , then?

$$\frac{\partial}{\partial u} \left( \frac{h^{-1}(u)}{1 - h^{-1}(u) \left( 1 - F^\alpha \left( \frac{x - \lambda}{\theta} \right) \right)} \right) \stackrel{\text{sign}}{=} \frac{\partial h^{-1}(u)}{\partial u} \leq 0,$$

implying that  $\frac{h^{-1}(u)}{1-h^{-1}(u)(1-F^\alpha(\frac{x-\lambda}{\theta}))}$  is decreasing in  $u$ . Again,

$$\frac{\partial}{\partial \lambda} \left( \frac{F^\alpha \left( \frac{x-\lambda}{\theta} \right)}{1-h^{-1}(u)(1-F^\alpha \left( \frac{x-\lambda}{\theta} \right))} \right) \stackrel{\text{sign}}{=} - \left( \frac{\alpha}{\theta} \right) \tilde{r} \left( \frac{x-\lambda}{\theta} \right) F^\alpha \left( \frac{x-\lambda}{\theta} \right) (1-h^{-1}(u)) \leq 0,$$

implying that  $\frac{F^\alpha \left( \frac{x-\lambda}{\theta} \right)}{1-h^{-1}(u)(1-F^\alpha \left( \frac{x-\lambda}{\theta} \right))}$  is decreasing in  $\lambda$ . Also,

$$\begin{aligned} \frac{\partial}{\partial \theta} \left( \frac{F^\alpha \left( \frac{x-\lambda}{\theta} \right)}{1-h^{-1}(u)(1-F^\alpha \left( \frac{x-\lambda}{\theta} \right))} \right) &\stackrel{\text{sign}}{=} - \left( \frac{\alpha}{\theta^2} \right) (x-\lambda) \tilde{r} \left( \frac{x-\lambda}{\theta} \right) F^\alpha \left( \frac{x-\lambda}{\theta} \right) \\ &\times (1-h^{-1}(u)) \leq 0, \end{aligned}$$

implying that  $\frac{F^\alpha \left( \frac{x-\lambda}{\theta} \right)}{1-h^{-1}(u)(1-F^\alpha \left( \frac{x-\lambda}{\theta} \right))}$  is decreasing in  $\theta$ . Thus, as  $\alpha, \beta, h(p), \lambda, \theta, \in \varepsilon_+$ , for  $i \leq j$  taking  $\alpha_i \leq \alpha_j$ ,  $u_i \leq u_j$ ,  $\lambda_i \leq \lambda_j$ ,  $\theta_i \leq \theta_j$  and noticing that  $h^{-1}(u)$  is decreasing in  $u$ , it can be written that

$$\begin{aligned} \frac{h^{-1}(u_j) F^{\alpha_j} \left( \frac{x-\lambda_j}{\theta_j} \right)}{1-h^{-1}(u_j) \left( 1-F^{\alpha_j} \left( \frac{x-\lambda_j}{\theta_j} \right) \right)} &\leq \frac{h^{-1}(u_j) F^{\alpha_i} \left( \frac{x-\lambda_j}{\theta_j} \right)}{1-h^{-1}(u_j) \left( 1-F^{\alpha_i} \left( \frac{x-\lambda_j}{\theta_j} \right) \right)} \\ &\leq \frac{h^{-1}(u_i) F^{\alpha_i} \left( \frac{x-\lambda_j}{\theta_j} \right)}{1-h^{-1}(u_i) \left( 1-F^{\alpha_i} \left( \frac{x-\lambda_j}{\theta_j} \right) \right)} \leq \frac{h^{-1}(u_i) F^{\alpha_i} \left( \frac{x-\lambda_i}{\theta_j} \right)}{1-h^{-1}(u_i) \left( 1-F^{\alpha_i} \left( \frac{x-\lambda_i}{\theta_j} \right) \right)} \\ &\leq \frac{h^{-1}(u_i) F^{\alpha_i} \left( \frac{x-\lambda_i}{\theta_i} \right)}{1-h^{-1}(u_i) \left( 1-F^{\alpha_i} \left( \frac{x-\lambda_i}{\theta_i} \right) \right)}, \end{aligned}$$

Which implies

$$\frac{h^{-1}(u_j) F^{\alpha_j} \left( \frac{x-\lambda_j}{\theta_j} \right)}{1-h^{-1}(u_j) \left( 1-F^{\alpha_j} \left( \frac{x-\lambda_j}{\theta_j} \right) \right)} \leq \frac{h^{-1}(u_i) F^{\alpha_i} \left( \frac{x-\lambda_i}{\theta_i} \right)}{1-h^{-1}(u_i) \left( 1-F^{\alpha_i} \left( \frac{x-\lambda_i}{\theta_i} \right) \right)},$$

Hence substituting the above results, we get  $\frac{\partial F_{Y_{n:n}}(x)}{\partial \alpha_i} - \frac{\partial F_{Y_{n:n}}(x)}{\partial \alpha_j} \leq 0$ . Thus by Lemma 3.3 of Kundu et al. (2016), it can be proved that  $F_{Y_{n:n}}(x)$  is Schur-convex in  $\alpha$ . thus the result is proved by Theorem A.8 of Marshall et al.(2011)  $\square$

**Theorem 3.6.** *Let Assumption 3.1 hold and  $h : [0, 1] \rightarrow R$  be a differentiable function. Also,  $\lambda_i = \lambda_i^* = \lambda$ ,  $\theta_i = \theta_i^* = \theta$ ,  $\alpha_i^* = \beta_i$ , where  $i = 1, \dots, n$ . Then,*

- (i)  $\alpha \succsim_w \beta$  implies  $Y_{1:n} \leq_{st} Y_{1:n}^*$ , provided  $\alpha, \beta \in \varepsilon_+$ ,  $\prod_{i=1}^n p_i \leq \prod_{i=1}^n p_i^*$ .
- (ii)  $\alpha^{-1} \succsim_w \beta^{-1}$  implies  $Y_{1:n} \leq_{st} Y_{1:n}^*$ , provided  $\alpha, \beta \in \varepsilon_+$ ,  $\prod_{i=1}^n p_i \leq \prod_{i=1}^n p_i^*$ .

**Proof(i)** : The survival function of  $Y_{1:n}$  is given by

$$\bar{F}_{Y_{1:n}}(x) = \prod_{i=1}^n p_i \left[ 1 - F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right) \right],$$

where  $h(p_i) = u_i$ . To prove the required result, it sufficient to show that  $\bar{F}_{Y_{1:n}}(x) \leq \bar{F}_{Y_{1:n}^*}(x)$ , which is equivalent to proving that  $\prod_{i=1}^n \left[ 1 - F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right) \right] \leq \prod_{i=1}^n \left[ 1 - F^{\beta_i} \left( \frac{x-\lambda}{\theta} \right) \right]$ , since  $\prod_{i=1}^n p_i \leq \prod_{i=1}^n p_i^*$ . Let  $\phi(\alpha) = \prod_{i=1}^n \left[ 1 - F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right) \right]$ . Differentiating  $\phi(\alpha)$ , partially, with respect to  $\alpha_i$ , we get

$$\frac{\partial \phi(\alpha)}{\partial \alpha_i} = - \frac{F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right) \ln[F \left( \frac{x-\lambda}{\theta} \right)]}{1 - F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right)} \phi(\alpha) \geq 0,$$

so,  $\phi(\alpha)$  is increasing in each  $\alpha_i$ . Again, it can be shown that

$$(3.5) \quad \frac{\partial \phi(\alpha)}{\partial \alpha_i} - \frac{\partial \phi(\alpha)}{\partial \alpha_j} \stackrel{\text{sign}}{\leq} \frac{F^{\alpha_j} \left( \frac{x-\lambda}{\theta} \right) \ln[F \left( \frac{x-\lambda}{\theta} \right)]}{1 - F^{\alpha_j} \left( \frac{x-\lambda}{\theta} \right)} - \frac{F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right) \ln[F \left( \frac{x-\lambda}{\theta} \right)]}{1 - F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right)}$$

Now,

$$\frac{\partial}{\partial \alpha} \left( \frac{F^\alpha \left( \frac{x-\lambda}{\theta} \right)}{1 - F^\alpha \left( \frac{x-\lambda}{\theta} \right)} \right) \stackrel{\text{sign}}{\leq} F^\alpha \left( \frac{x-\lambda}{\theta} \right) \ln[F \left( \frac{x-\lambda}{\theta} \right)] \leq 0,$$

implying that  $\frac{F^\alpha \left( \frac{x-\lambda}{\theta} \right)}{1 - F^\alpha \left( \frac{x-\lambda}{\theta} \right)}$  is decreasing in  $\alpha$ . Thus, as  $\alpha \in \varepsilon_+$ , for  $i \leq j$  taking  $\alpha_i \leq \alpha_j$ , it can be written that

$$\frac{F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right)}{1 - F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right)} \geq \frac{F^{\alpha_j} \left( \frac{x-\lambda}{\theta} \right)}{1 - F^{\alpha_j} \left( \frac{x-\lambda}{\theta} \right)},$$

which implies

$$\frac{F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right) \ln[F \left( \frac{x-\lambda}{\theta} \right)]}{1 - F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right)} \leq \frac{F^{\alpha_j} \left( \frac{x-\lambda}{\theta} \right) \ln[F \left( \frac{x-\lambda}{\theta} \right)]}{1 - F^{\alpha_j} \left( \frac{x-\lambda}{\theta} \right)},$$

Hence, from (3.5), we get  $\frac{\partial \phi(\alpha)}{\partial \alpha_i} - \frac{\partial \phi(\alpha)}{\partial \alpha_j} \geq 0$ . Thus, by Lemma 3.3 of Kundu et al. (2016), it can be proved that  $\phi(\alpha)$  is Schur-concave in  $\alpha$ . Thus the result is proved by Theorem A.8 of Marshall et al. (2011).

**Proof(ii)** : The survival function of  $Y_{1:n}$  can be expressed as the function of  $c_i = \frac{1}{\alpha_i}$ ,  $i = 1, \dots, n$ . We denote it by  $\psi_5(\zeta)$ , where  $\zeta = (c_1, \dots, c_n)$ , and

$$\psi_5(\alpha) = \prod_{i=1}^n p_i \left[ 1 - F^{\frac{1}{c_i}} \left( \frac{x-\lambda}{\theta} \right) \right],$$

where  $h(p_i) = u_i$ . To prove the required result, it is sufficient to show that  $\psi_5(\underline{\alpha}) \leq \psi_5(\underline{\beta})$ , which is equivalent to proving that  $\prod_{i=1}^n \left[ 1 - F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right) \right] \leq \prod_{i=1}^n \left[ 1 - F^{\beta_i} \left( \frac{x-\lambda}{\theta} \right) \right]$ , since  $\prod_{i=1}^n p_i \leq \prod_{i=1}^n p_i^*$ . Let  $\phi(\underline{c}) = \prod_{i=1}^n \left[ 1 - F^{\frac{1}{c_i}} \left( \frac{x-\lambda}{\theta} \right) \right]$ . It can be shown that the partial derivative of  $\phi(\underline{c})$ , with respect to  $c_i$ , is decreasing in each  $c_i$ . Thus, by Lemma 3.1 of Kundu et al. (2016), it can be proved that  $\phi(\underline{c})$  is Schur-concave in  $\underline{c}$ . Thus the result is proved by Theorem A.8 of Marshall et al. (2011).

**Theorem 3.7.** *Let Assumption 3.1 hold and  $h : [0, 1] \rightarrow R$  be a differentiable function. Also,  $\lambda_i = \lambda_i^* = \lambda$ ,  $\theta_i = \theta_i^* = \theta$ , and  $\alpha_i^* = \beta_i$ , where  $i = 1, \dots, n$ . Then,  $\underline{\alpha} \underset{\sim}{\geq}^p \underline{\beta}$  implies  $Y_{1:n} \leq_{st} Y_{1:n}^*$ , provided  $\underline{\alpha}, \underline{\beta} \in D_+$ ,  $\prod_{i=1}^n p_i \leq \prod_{i=1}^n p_i^*$ .*

**Proof:** The survival function of  $Y_{1:n}$  can be expressed as the function of  $d_i$ , where  $d_i = \ln \alpha_i$ ,  $i = 1, \dots, n$ . We denote it by  $\psi_6(\underline{d})$ , where  $\underline{d} = (d_1, \dots, d_n)$ .

$$\psi_6(\underline{d}) = \prod_{i=1}^n p_i \left[ 1 - F^{e^{d_i}} \left( \frac{x-\lambda}{\theta} \right) \right],$$

where  $h(p_i) = u_i$ . To prove the required result, it is sufficient to show that  $\psi_6(\underline{\alpha}) \leq \psi_6(\underline{\beta})$ , which is equivalent to proving that  $\prod_{i=1}^n p_i \left[ 1 - F^{\alpha_i} \left( \frac{x-\lambda}{\theta} \right) \right] \leq \prod_{i=1}^n p_i \left[ 1 - F^{\beta_i} \left( \frac{x-\lambda}{\theta} \right) \right]$ , since  $\prod_{i=1}^n p_i \leq \prod_{i=1}^n p_i^*$ . Let  $\phi(\underline{d}) = \prod_{i=1}^n \left[ 1 - F^{e^{d_i}} \left( \frac{x-\lambda}{\theta} \right) \right]$ . Differentiating  $\psi_6(\underline{d})$ , partially, with respect to  $d_i$ , we get

$$\frac{\partial \phi(\underline{d})}{\partial d_i} = - \frac{e^{d_i} F^{e^{d_i}} \left( \frac{x-\lambda}{\theta} \right) \ln[F \left( \frac{x-\lambda}{\theta} \right)]}{1 - F^{e^{d_i}} \left( \frac{x-\lambda}{\theta} \right)} \phi(\underline{d}) \geq 0$$

so,  $\phi(\underline{d})$  is increasing in each  $d_i$ . Again, it can be shown that

$$(3.6) \quad \frac{\partial \phi(\underline{d})}{\partial d_i} - \frac{\partial \phi(\underline{d})}{\partial d_j} \underset{\text{sign}}{=} \frac{e^{d_j} F^{e^{d_j}} \left( \frac{x-\lambda}{\theta} \right) \ln[F \left( \frac{x-\lambda}{\theta} \right)]}{1 - F^{e^{d_j}} \left( \frac{x-\lambda}{\theta} \right)} - \frac{e^{d_i} F^{e^{d_i}} \left( \frac{x-\lambda}{\theta} \right) \ln[F \left( \frac{x-\lambda}{\theta} \right)]}{1 - F^{e^{d_i}} \left( \frac{x-\lambda}{\theta} \right)}$$

Now,

$$\frac{\partial}{\partial d} \left( \frac{F^{e^d} \left( \frac{x-\lambda}{\theta} \right)}{1 - F^{e^d} \left( \frac{x-\lambda}{\theta} \right)} \right) \underset{\text{sign}}{=} e^d F^{e^d} \left( \frac{x-\lambda}{\theta} \right) \ln[F \left( \frac{x-\lambda}{\theta} \right)] \leq 0,$$

implying that  $\frac{F^{e^d} \left( \frac{x-\lambda}{\theta} \right)}{1 - F^{e^d} \left( \frac{x-\lambda}{\theta} \right)}$  is decreasing in  $d$ . Thus, as  $\underline{d} \in D_+$ , for  $i \leq j$  taking  $d_i \geq d_j$ , it can be written that

$$\frac{F^{e^{d_i}} \left( \frac{x-\lambda}{\theta} \right)}{1 - F^{e^{d_i}} \left( \frac{x-\lambda}{\theta} \right)} \leq \frac{F^{e^{d_j}} \left( \frac{x-\lambda}{\theta} \right)}{1 - F^{e^{d_j}} \left( \frac{x-\lambda}{\theta} \right)},$$

which implies

$$\frac{e^{d_i} F^{e^{d_i}} \left( \frac{x-\lambda}{\theta} \right) \ln[F \left( \frac{x-\lambda}{\theta} \right)]}{1 - F^{e^{d_i}} \left( \frac{x-\lambda}{\theta} \right)} \geq \frac{e^{d_j} F^{e^{d_j}} \left( \frac{x-\lambda}{\theta} \right) \ln[F \left( \frac{x-\lambda}{\theta} \right)]}{1 - F^{e^{d_j}} \left( \frac{x-\lambda}{\theta} \right)}.$$

Hence, from (3.6), we get  $\frac{\partial \phi(\underline{d})}{\partial d_i} - \frac{\partial \phi(\underline{d})}{\partial d_j} \leq 0$ . Thus, by Lemma 3.1 of Kundu et al. (2016), it can be proved that  $\phi(\underline{d})$  is Schur-concave in  $\underline{d}$ . Thus the result is proved by Lemma 3.1 of Khaledi et al. (2002).  $\square$

**Theorem 3.8.** *Let Assumption 3.1 hold and  $h : [0, 1] \rightarrow R$  be a differentiable function. Also,  $\lambda_i = \lambda_i^* = \lambda$ ,  $\theta_i^* = \delta_i$ , and  $\alpha_i = \alpha_i^* = \alpha$ , where  $i = 1, \dots, n$ . Then,  $\varrho^{-1} \geq_w \varrho^{-1}$  implies  $Y_{1:n} \leq_{st} Y_{1:n}^*$ , provided  $\vartheta, \vardelta \in \varepsilon_+$ ,  $\prod_{i=1}^n p_i \leq \prod_{i=1}^n p_i^*$ , and  $\tilde{r}(u)$  is increasing in  $u$ .*

**Proof:** The survival function of  $Y_{1:n}$  can be expressed as the function of  $e_i$ , where  $e_i = \frac{1}{\theta_i}$ ,  $i = 1, \dots, n$ . We denote it by  $\psi_7(\underline{e})$ , where  $\underline{e} = (e_1, \dots, e_n)$ , and

$$\psi_7(\underline{e}) = \prod_{i=1}^n p_i \left[ 1 - F^\alpha(e_i(x - \lambda)) \right],$$

where  $h(p_i) = u_i$ . To prove the required result, it is sufficient to show that  $\psi_7(\varrho^{-1}) \leq \psi_7(\vardelta^{-1})$ , which is equivalent to proving that  $\prod_{i=1}^n \left[ 1 - F^\alpha\left(\frac{1}{\theta_i}(x - \lambda)\right) \right] \leq \prod_{i=1}^n \left[ 1 - F^\alpha\left(\frac{1}{\delta_i}(x - \lambda)\right) \right]$ , since  $\prod_{i=1}^n p_i \leq \prod_{i=1}^n p_i^*$ . Let  $\phi(\underline{e}) = \prod_{i=1}^n \left[ 1 - F^\alpha(e_i(x - \lambda)) \right]$ . Differentiating  $\phi(\underline{e})$ , partially, with respect to  $e_i$ , we get

$$\frac{\partial \phi(\underline{e})}{\partial e_i} = - \frac{\alpha(x - \lambda) \tilde{r}(e_i(x - \lambda)) F^\alpha(e_i(x - \lambda))}{1 - F^\alpha(e_i(x - \lambda))} \phi(\underline{e}) \leq 0,$$

so,  $\psi_7(\underline{e})$  is decreasing in each  $e_i$ . Again, it can be shown that

$$(3.7) \quad \frac{\partial \phi(\underline{e})}{\partial e_i} - \frac{\partial \phi(\underline{e})}{\partial e_j} \stackrel{\text{sign}}{=} \frac{\alpha(x - \lambda) \tilde{r}(e_j(x - \lambda)) F^\alpha(e_j(x - \lambda))}{1 - F^\alpha(e_j(x - \lambda))} - \frac{\alpha(x - \lambda) \tilde{r}(e_i(x - \lambda)) F^\alpha(e_i(x - \lambda))}{1 - F^\alpha(e_i(x - \lambda))}.$$

Now,

$$\frac{\partial}{\partial e} \left( \frac{F^\alpha(e(x - \lambda))}{1 - F^\alpha(e(x - \lambda))} \right) \stackrel{\text{sign}}{=} \alpha(x - \lambda) \tilde{r}(e(x - \lambda)) F^\alpha(e(x - \lambda)) \geq 0,$$

implying that  $\frac{F^\alpha(e(x - \lambda))}{1 - F^\alpha(e(x - \lambda))}$  is increasing in  $e$ . Thus, us  $\vartheta \in \varepsilon_+$ , for  $i \leq j$  taking  $e_i \geq e_j$ , it can be written that

$$\frac{F^\alpha(e_i(x - \lambda))}{1 - F^\alpha(e_i(x - \lambda))} \geq \frac{F^\alpha(e_j(x - \lambda))}{1 - F^\alpha(e_j(x - \lambda))},$$



As  $\tilde{r}(u)$  is increasing in  $u$ , then

$$\frac{\alpha(x-\lambda)\tilde{r}(e_i(x-\lambda))F^\alpha(e_i(x-\lambda))}{1-F^\alpha(e_i(x-\lambda))} \geq \frac{\alpha(x-\lambda)\tilde{r}(e_j(x-\lambda))F^\alpha(e_j(x-\lambda))}{1-F^\alpha(e_j(x-\lambda))}$$

Hence, from (3.7), we get  $\frac{\partial\phi(\underline{\epsilon})}{\partial e_i} - \frac{\partial\phi(\underline{\epsilon})}{\partial e_j} \leq 0$ . Thus, by Lemma 3.1 of Kundu et al. (2016), it can be proved that  $\phi(\underline{\epsilon})$  is Schur-concave in  $\underline{\epsilon}$ . Thus the result is proved by Theorem A.8 Marshall et al. (2011).  $\square$

Theorem 3.8 shows that if both the location and shock parameter vectors i.e. the matrix of location and shock parameters of one system majorizes the other when the scale and shape parameter vectors remain constant do not lead to better system reliability.

**Theorem 3.9.** For  $n = 2$ , let Assumption 3.1 hold. Further, let  $h : [0, 1] \rightarrow R_+$  be a differentiable and strictly increasing concave function. Then, for  $i = 1, 2$ , if  $\theta_i = \theta_i^* = \alpha_i = \alpha_i^* = \theta$ , and  $(h(p), \lambda) \in P_2$ , we have that

$$\begin{pmatrix} h(p_1) & h(p_2) \\ \lambda_1 & \lambda_2 \end{pmatrix} \gg \begin{pmatrix} h(p_1^*) & h(p_2^*) \\ \lambda_1^* & \lambda_2^* \end{pmatrix},$$

implies  $Y_{1:2}^* \geq_{st} Y_{1:2}$ , provided  $\tilde{r}(u)$  is increasing in  $u$ .

**Proof :** With  $u_1 = h(p_1)$ ,  $u_2 = h(p_2)$ , we have  $p_1 = h^{-1}(u_1)$ ,  $p_2 = h^{-1}(u_2)$ , where  $h^{-1}$  denotes the inverse of the function  $h$ . From (1.2), the survival function of  $Y_{1:2}$  is,

$$\bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \theta, \theta) = \prod_{i=1}^2 h^{-1}(u_i) \left[ 1 - F^\theta \left( \frac{x - \lambda_i}{\theta} \right) \right], \quad x > \max\{\lambda_i, i = 1, \dots, n\}.$$

Note that the function  $\bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \theta, \theta)$  is permutation invariant in  $(u_i, \lambda_i)$ , and so condition (i) of Theorem 2 of Balakrishnan et al. (2015) is satisfied. Next, we have to show that condition (ii) of Theorem 2 of Balakrishnan et al. (2015) also holds. The assumption  $(\underline{y}, \underline{\lambda}) \in P_2$  implies that  $(u_1 - u_2)(\lambda_1 - \lambda_2) \leq 0$ . This implies that  $u_1 \geq u_2$  and  $\lambda_1 \leq \lambda_2$  or  $u_1 \leq u_2$  and  $\lambda_1 \geq \lambda_2$ . We proof only for the case when  $u_1 \geq u_2$  and  $\lambda_1 \leq \lambda_2$ . The proof for the other case is similar. The partial derivatives of  $\bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \theta, \theta)$  with respect to  $u_i$  and  $\lambda_i$  are

$$\begin{aligned} \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \theta, \theta)}{\partial u_i} &= \frac{\partial h^{-1}(u_i)}{\partial u_i} \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \theta, \theta), \\ \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \theta, \theta)}{\partial \lambda_i} &= - \frac{\tilde{r} \left( \frac{x - \lambda_i}{\theta} \right) F^\theta \left( \frac{x - \lambda_i}{\theta} \right)}{1 - F^\theta \left( \frac{x - \lambda_i}{\theta} \right)} \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \theta, \theta). \end{aligned}$$

For fixed  $x > \max\{\lambda_i, i = 1, \dots, n\}$ , let us define the function  $\varphi(\underline{y}, \underline{\lambda})$  as follows:

$$\begin{aligned}
(3.8) \quad \varphi(\underline{y}, \underline{\lambda}) &= (u_1 - u_2) \left( \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \theta, \theta)}{\partial u_1} - \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \theta, \theta)}{\partial u_2} \right) \\
&+ (\lambda_1 - \lambda_2) \left( \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \theta, \theta)}{\partial \lambda_1} - \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \theta, \theta)}{\partial \lambda_2} \right) \\
&= (u_1 - u_2) \left( \frac{\frac{\partial h^{-1}(u_1)}{\partial u_1}}{h^{-1}(u_1)} - \frac{\frac{\partial h^{-1}(u_2)}{\partial u_2}}{h^{-1}(u_2)} \right) \times \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \theta, \theta) \\
&+ (\lambda_1 - \lambda_2) \left( \frac{\tilde{r}\left(\frac{x-\lambda_1}{\theta}\right) F^\theta\left(\frac{x-\lambda_1}{\theta}\right)}{1 - F^\theta\left(\frac{x-\lambda_1}{\theta}\right)} - \frac{\tilde{r}\left(\frac{x-\lambda_2}{\theta}\right) F^\theta\left(\frac{x-\lambda_2}{\theta}\right)}{1 - F^\theta\left(\frac{x-\lambda_2}{\theta}\right)} \right) \\
&\times \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \theta, \theta).
\end{aligned}$$

Since  $h$  is strictly increasing and concave, then for  $u_1 \geq u_2$  and  $\lambda_1 \leq \lambda_2$ , we have

$$(3.9) \quad \frac{\frac{\partial h^{-1}(u_1)}{\partial u_1}}{h^{-1}(u_1)} \leq \frac{\frac{\partial h^{-1}(u_2)}{\partial u_2}}{h^{-1}(u_2)}.$$

Furthermore,  $\frac{t^\theta}{1-t^\theta}$  is increasing in  $t$  for  $\theta > 0$ . For the reversed hazard rate function  $\tilde{r}$  that is increasing, we have

$$(3.10) \quad \tilde{r}\left(\frac{x-\lambda_1}{\theta}\right) \geq \tilde{r}\left(\frac{x-\lambda_2}{\theta}\right),$$

and

$$(3.11) \quad \frac{\tilde{r}\left(\frac{x-\lambda_1}{\theta}\right) F^\theta\left(\frac{x-\lambda_1}{\theta}\right)}{1 - F^\theta\left(\frac{x-\lambda_1}{\theta}\right)} \geq \frac{\tilde{r}\left(\frac{x-\lambda_2}{\theta}\right) F^\theta\left(\frac{x-\lambda_2}{\theta}\right)}{1 - F^\theta\left(\frac{x-\lambda_2}{\theta}\right)}.$$

Combining (3.9) and (3.10) and (3.11), we see that  $\varphi(\underline{y}, \underline{\lambda}) \leq 0$ . Condition (ii) Theorem 2 of Balakrishnan et al. (2015) is satisfied, which completes the proof.

**Counterexample 3.1** Let the baseline distribution function be  $F(t) = e^{-\frac{1}{t}}$ ,  $t > 0$ . Take  $h(p) = -\ln p$ . Here, the baseline reversed hazard rate function is decreasing and  $h(p)$  is decreasing and convex. Thus, the assumptions of the Theorem 3.9 are violated. Let us take  $\theta_1 = \theta_2 = \theta_1^* = \theta_2^* = \alpha_1 = \alpha_2 = \alpha_1^* = \alpha_2^* = 1.5$ ,  $(\lambda_1, \lambda_2) = (0.3, 0.9)$ ,  $(\lambda_1^*, \lambda_2^*) = (0.54, 0.66)$ ,  $(p_1, p_2) = (e^{-0.4}, e^{-0.5})$  and  $(p_1^*, p_2^*) = (e^{-0.44}, e^{-0.46})$ . Consider the T-transform matrix  $T_{0.6} = \begin{pmatrix} 0.6 & 0.4 \\ 0.4 & 0.6 \end{pmatrix}$ .

It can be shown that

$$\begin{pmatrix} h(p_1^*) & h(p_2^*) \\ \lambda_1^* & \lambda_2^* \end{pmatrix} = \begin{pmatrix} h(p_1) & h(p_2) \\ \lambda_1 & \lambda_2 \end{pmatrix} T_{0.6},$$

which implies  $\begin{pmatrix} h(p_1) & h(p_2) \\ \lambda_1 & \lambda_2 \end{pmatrix} \gg \begin{pmatrix} h(p_1^*) & h(p_2^*) \\ \lambda_1^* & \lambda_2^* \end{pmatrix}$ . Under this set up,  $\bar{F}_{Y_{1:2}}(2) = 0.2597610428$ ,  $\bar{F}_{Y_{1:2}^*}(2) = 0.2599036428$ ,  $\bar{F}_{Y_{1:2}}(5) = 0.06532417018$ ,  $\bar{F}_{Y_{1:2}^*}(5) = 0.06516286182$ , which readily shows that  $Y_{1:2}^* \not\geq_{st} Y_{1:2}$ .

The following theorem extends Theorem 3.8 when two sets of  $n$ -independent observations from  $ELS$  distribution. The generalization is the direct result of the Theorem 3.8 and Lemma 5 of Balakrishnan et al. (2018). So, the proof is omitted.

**Theorem 3.10.** *Let Assumption 3.1 hold and  $h : [0, 1] \rightarrow R_+$  be a differentiable and strictly increasing concave function. Further, let  $T_w$  be a T-transform matrix. Then, for  $i = 1, \dots, n$ , if  $\theta_i = \theta_i^* = \alpha_i = \alpha_i^* = \theta$ , and  $(h(p), \lambda) \in P_n$ , we have that*

$$\begin{pmatrix} h(p_1^*) & h(p_2^*) & \cdots & h(p_n^*) \\ \lambda_1^* & \lambda_2^* & \cdots & \lambda_n^* \end{pmatrix} = \begin{pmatrix} h(p_1) & h(p_2) & \cdots & h(p_n) \\ \lambda_1 & \lambda_2 & \cdots & \lambda_n \end{pmatrix} T_w,$$

implies  $Y_{1:n}^* \geq_{st} Y_{1:n}$ , provided  $\tilde{r}(u)$  is increasing in  $u$ .

**Theorem 3.11.** *Let Assumption 3.1 hold. Further, let  $T_{w_1}, \dots, T_{w_k}$  have the same structure. Suppose  $h : [0, 1] \rightarrow R_+$  is a differentiable and strictly increasing concave function. Then, for  $i = 1, \dots, n$ , if  $\theta_i = \theta_i^* = \alpha_i = \alpha_i^* = \theta$ , and  $(h(p), \lambda) \in P_n$ , we have that*

$$\begin{pmatrix} h(p_1^*) & h(p_2^*) & \cdots & h(p_n^*) \\ \lambda_1^* & \lambda_2^* & \cdots & \lambda_n^* \end{pmatrix} = \begin{pmatrix} h(p_1) & h(p_2) & \cdots & h(p_n) \\ \lambda_1 & \lambda_2 & \cdots & \lambda_n \end{pmatrix} T_{w_1} \cdots T_{w_k},$$

implies  $Y_{1:n}^* \geq_{st} Y_{1:n}$ , provided  $\tilde{r}(u)$  is increasing in  $u$ .

**Proof:** Since a finite product of T-transform matrices with the same structure is also a T-transform matrix, so, the desired result is obtained from Theorem 3.9.  $\square$

Our next Theorem shows that the result in Theorem 3.10 holds for T-transform matrices with different structure.

**Theorem 3.12.** *Let Assumption 3.1 hold. Further, let  $T_{w_1}, \dots, T_{w_k}, k > 2$ , have different structures. Suppose  $h : [0, 1] \rightarrow R_+$  is a differentiable and strictly increasing concave function. Then, for  $i = 1, \dots, n$ , if  $\theta_i = \theta_i^* = \alpha_i = \alpha_i^* = \theta$ ,  $(h(p), \lambda) \in P_n$  and  $(h(p), \lambda) T_{w_1} \cdots T_{w_k} \in P_n$ , we have that*

$$\begin{pmatrix} h(p_1^*) & h(p_2^*) & \cdots & h(p_n^*) \\ \lambda_1^* & \lambda_2^* & \cdots & \lambda_n^* \end{pmatrix} = \begin{pmatrix} h(p_1) & h(p_2) & \cdots & h(p_n) \\ \lambda_1 & \lambda_2 & \cdots & \lambda_n \end{pmatrix} T_{w_1} \cdots T_{w_k},$$

implies  $Y_{1:n}^* \geq_{st} Y_{1:n}$ , provided  $\tilde{r}(u)$  is increasing in  $u$ .

**Proof:**

$$\begin{pmatrix} h(p_1)^{(i)} & \cdots & h(p_n)^{(i)} \\ \lambda_1^{(i)} & \cdots & \lambda_n^{(i)} \end{pmatrix} = \begin{pmatrix} h(p_1) & \cdots & h(p_n) \\ \lambda_1 & \cdots & \lambda_n \end{pmatrix} T_{w_1} \cdots T_{w_i}, \text{ for } i = 1, \dots, k.$$

Assume  $V_1^{(i)}, \dots, V_n^{(i)}$ ,  $i = 1, \dots, k$ , are independent sets of random variables with

$V_j^{(i)} \sim ELS(\lambda_j^{(i)}, \theta_j, \alpha_j)$  where  $\theta_j = \theta_j^* = \alpha_j = \alpha_j^* = \theta$ ,  $j = 1, \dots, n$  and  $i = 1, \dots, k$ . From the assumption of the theorem, it follows that

$$\begin{pmatrix} h(p_1)^{(i)} & \cdots & h(p_n)^{(i)} \\ \lambda_1^{(i)} & \cdots & \lambda_n^{(i)} \end{pmatrix} \in P_n, \text{ for } i = 1, \dots, k.$$

From these observations and the results of Theorem 3.9, it then follows that

$$Y_{1:n} \leq_{st} V_{1:n}^{(1)} \leq_{st} \cdots \leq_{st} V_{1:n}^{(k-2)} \leq_{st} V_{1:n}^{(k-1)} \leq_{st} Y_{1:n}^*.$$

Which completes the proof of the theorem.  $\square$

The following example illustrates the result established in Theorem 3.11.

**Example 3.1** Suppose  $X_1, X_2$  and  $X_3$  are independent non-negative random variables with  $X_i \sim ELS(\lambda_i, \theta_i, \alpha_i)$ , and  $I_{p_1}, I_{p_2}$  and  $I_{p_3}$  are independent Bernoulli random variables, independent of  $X_i$ 's, with  $E(I_{p_i}) = p_i$ ,  $i = 1, 2, 3$ . Further, suppose  $X_1^*, X_2^*$  and  $X_3^*$  are independent non-negative random variables with  $X_i^* \sim ELS(\lambda_i^*, \theta_i^*, \alpha_i^*)$ , and  $I_{p_1^*}, I_{p_2^*}$  and  $I_{p_3^*}$  are independent Bernoulli random variables, independent of  $X_i^*$ 's, with  $E(I_{p_i^*}) = p_i^*$ ,  $i = 1, 2, 3$ . Consider a baseline distribution with distribution function  $F(x) = 1 - e^{-x}$ ,  $x > 0$ . Consider the T-transform matrices as follows:

$$T_{0.7} = \begin{pmatrix} 0.7 & 0 & 0.3 \\ 0 & 1 & 0 \\ 0.3 & 0 & 0.7 \end{pmatrix}, T_{0.6} = \begin{pmatrix} 0.4 & 0.6 & 0 \\ 0.6 & 0.4 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

$$T_{0.4} = \begin{pmatrix} 0.6 & 0 & 0.4 \\ 0 & 1 & 0 \\ 0.4 & 0 & 0.6 \end{pmatrix}.$$

Suppose  $h(p) = \frac{p}{1+p}$ . Then, for  $\theta_i = \theta_i^* = \alpha_i = \alpha_i^* = 1$ ,  $i = 1, 2, 3$ , let  $(\lambda_1, \lambda_2, \lambda_3) = (2, 3, 4)$ ,  $(\lambda_1^*, \lambda_2^*, \lambda_3^*) = (3.06, 2.76, 3.17)$ ,  $(p_1, p_2, p_3) = (0.4, 0.3, 0.2)$  and  $(p_1^*, p_2^*, p_3^*) = (0.285, 0.317, 0.273)$ . It is easy to observe that  $(h(p), \lambda) \in P_3$ ,  $(h(p), \lambda)T_{0.7} \in P_3$  and  $(h(p), \lambda)T_{0.7}T_{0.6} \in P_3$  and  $(h(p), \lambda)T_{0.7}T_{0.6}T_{0.4} \in P_3$ .

So, from Theorem 3.11, we have  $Y_{1:3}^* \geq_{st} Y_{1:3}$ .

**Theorem 3.13.** *Let Assumption 3.1 hold for  $n = 2$ . Suppose  $h : [0, 1] \rightarrow R_+$  is differentiable and strictly increasing concave function. Further, let  $\tilde{r}(u)$  and  $u\tilde{r}(u)$  are increasing in  $u$ . Then, if  $\alpha_i = \alpha_i^* = \alpha$ , and  $(h(p), \lambda, \varrho) \in S_2$ , we have that*

$$\begin{pmatrix} h(p_1) & h(p_2) \\ \lambda_1 & \lambda_2 \\ \theta_1 & \theta_2 \end{pmatrix} \gg \begin{pmatrix} h(p_1^*) & h(p_2^*) \\ \lambda_1^* & \lambda_2^* \\ \theta_1^* & \theta_2^* \end{pmatrix},$$

implies  $Y_{1:2}^* \geq_{st} Y_{1:2}$ .

**Proof:** With  $u_1 = h(p_1), u_2 = h(p_2)$ , we have  $p_1 = h^{-1}(u_1), p_2 = h^{-1}(u_2)$ , where  $h^{-1}$  denotes the inverse of the function  $h$ . From (1.2), the survival function of  $Y_{1:2}$  is,

$$\bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha) = \prod_{i=1}^2 h^{-1}(u_i) \left[ 1 - F^\alpha \left( \frac{x - \lambda_i}{\theta_i} \right) \right], x > \max\{\lambda_i, i = 1, \dots, n\}.$$

Note that the function  $\bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha)$  is permutation invariant in  $(u_i, \lambda_i, \theta_i)$  and therefore condition (i) of Lemma 6 of Balakrishnan et al. (2018) is satisfied. Next, we have to show that Condition (ii) of Lemma 6 of Balakrishnan et al. (2018) also holds. The assumption  $(\underline{y}, \underline{\lambda}, \underline{\theta}) \in S_2$  implies that  $u_1 \leq (\geq) u_2$  and  $\lambda_1 \geq (\leq) \lambda_2$  and  $\theta_1 \geq (\leq) \theta_2$ . We proof only for the case when  $u_1 \leq u_2$  and  $\lambda_1 \geq \lambda_2$  and  $\theta_1 \geq \theta_2$ . The proof for the other case is similar. The partial derivatives of  $\bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha)$  with respect to  $u_i$  and  $\lambda_i$  and  $\theta_i$  are

$$\begin{aligned} \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha)}{\partial u_i} &= \frac{\partial h^{-1}(u_i)}{h^{-1}(u_i)} \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha), \\ \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha)}{\partial \lambda_i} &= \frac{\alpha}{\theta_i} \times \frac{\tilde{r} \left( \frac{x - \lambda_i}{\theta_i} \right) F^\alpha \left( \frac{x - \lambda_i}{\theta_i} \right)}{1 - F^\alpha \left( \frac{x - \lambda_i}{\theta_i} \right)} \times \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha), \\ \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha)}{\partial \theta_i} &= \frac{\alpha}{\theta_i} \times \frac{\left( \frac{x - \lambda_i}{\theta_i} \right) \tilde{r} \left( \frac{x - \lambda_i}{\theta_i} \right) F^\alpha \left( \frac{x - \lambda_i}{\theta_i} \right)}{1 - F^\alpha \left( \frac{x - \lambda_i}{\theta_i} \right)} \times \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha). \end{aligned}$$

For fixed  $x > \max\{\lambda_i, i = 1, \dots, n\}$ , let us define the function  $\varphi(\underline{y}, \underline{\lambda}, \underline{\theta})$  as follows:

$$\begin{aligned} \varphi(\underline{y}, \underline{\lambda}, \underline{\theta}) &= (u_1 - u_2) \left( \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha)}{\partial u_1} - \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha)}{\partial u_2} \right) \\ &+ (\lambda_1 - \lambda_2) \left( \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha)}{\partial \lambda_1} - \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha)}{\partial \lambda_2} \right) \\ &+ (\theta_1 - \theta_2) \left( \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha)}{\partial \theta_1} - \frac{\partial \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha)}{\partial \theta_2} \right) \\ &= (u_1 - u_2) \left( \frac{\partial h^{-1}(u_1)}{h^{-1}(u_1)} - \frac{\partial h^{-1}(u_2)}{h^{-1}(u_2)} \right) \times \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha) \\ &+ (\lambda_1 - \lambda_2) \left( \frac{1}{\theta_1} \tilde{r} \left( \frac{x - \lambda_1}{\theta_1} \right) l_1 - \frac{1}{\theta_2} \tilde{r} \left( \frac{x - \lambda_2}{\theta_2} \right) l_2 \right) \times \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha) \\ (3.12) \quad &+ (\theta_1 - \theta_2) \left( \frac{1}{\theta_1} \left( \frac{x - \lambda_1}{\theta_1} \right) \tilde{r} \left( \frac{x - \lambda_1}{\theta_1} \right) l_1 - \frac{1}{\theta_2} \left( \frac{x - \lambda_2}{\theta_2} \right) \tilde{r} \left( \frac{x - \lambda_2}{\theta_2} \right) l_2 \right) \\ &\times \bar{F}_{Y_{1:2}}(x; \underline{y}, \underline{\lambda}, \underline{\theta}, \alpha), \end{aligned}$$

where  $l_i = l(\alpha, F^\alpha \left( \frac{x - \lambda_i}{\theta_i} \right)) = \frac{\alpha F^\alpha \left( \frac{x - \lambda_i}{\theta_i} \right)}{1 - F^\alpha \left( \frac{x - \lambda_i}{\theta_i} \right)}$ ,  $i = 1, 2$ , is defined in Lemma 2.8 of Torrado (2015). Since  $h$  is strictly increasing and concave function, then for

$u_1 \leq u_2$  and  $\lambda_1 \geq \lambda_2$  and  $\theta_1 \geq \theta_2$ , we have

$$(3.13) \quad \frac{\frac{\partial h^{-1}(u_1)}{\partial u_1}}{h^{-1}(u_1)} \geq \frac{\frac{\partial h^{-1}(u_2)}{\partial u_2}}{h^{-1}(u_2)}.$$

$$(3.14) \quad \frac{1}{\theta_1} \tilde{r} \left( \frac{x - \lambda_1}{\theta_1} \right) l(\alpha, F^\alpha \left( \frac{x - \lambda_1}{\theta_1} \right)) \leq \frac{1}{\theta_2} \tilde{r} \left( \frac{x - \lambda_2}{\theta_2} \right) l(\alpha, F^\alpha \left( \frac{x - \lambda_2}{\theta_2} \right))$$

and

$$(3.15) \quad \frac{1}{\theta_1} \left( \frac{x - \lambda_1}{\theta_1} \right) \tilde{r} \left( \frac{x - \lambda_1}{\theta_1} \right) l(\alpha, F^\alpha \left( \frac{x - \lambda_1}{\theta_1} \right)) \leq \frac{1}{\theta_2} \left( \frac{x - \lambda_2}{\theta_2} \right) \tilde{r} \left( \frac{x - \lambda_2}{\theta_2} \right) l(\alpha, F^\alpha \left( \frac{x - \lambda_2}{\theta_2} \right))$$

combining (3.13), (3.14) and (3.15) in (3.12), we see that  $\varphi(y, \lambda, \theta) \leq 0$ . So condition (ii) of Lemma 6 of Balakrishnan et al. (2018) is satisfied, which completes the proof.  $\square$

**Counterexample 3.2** Let the baseline distribution function be  $F(t) = e^{-\frac{1}{t}}$ ,  $t > 0$ . Here,  $\tilde{r}(t)$  and  $t\tilde{r}(t)$  are decreasing. Take  $h(p) = p$ . Thus, the assumption of Theorem 3.13 are violated. Let us take  $\alpha_1 = \alpha_2 = \alpha_1^* = \alpha_2^* = 2.2$ ,  $(\theta_1, \theta_2) = (5.2, 2.7)$ ,  $(\theta_1^*, \theta_2^*) = (4.2, 3.7)$ ,  $(\lambda_1, \lambda_2) = (2.2, 2.5)$ ,  $(\lambda_1^*, \lambda_2^*) = (2.32, 2.38)$ ,  $(p_1, p_2) = (p_1^*, p_2^*) = (0.2, 0.2)$ . Consider the T-transform matrix  $T_{0.6} = \begin{pmatrix} 0.6 & 0.4 \\ 0.4 & 0.6 \end{pmatrix}$ . It can be shown that

$$\begin{pmatrix} h(p_1^*) & h(p_2^*) \\ \lambda_1^* & \lambda_2^* \\ \theta_1^* & \theta_2^* \end{pmatrix} = \begin{pmatrix} h(p_1) & h(p_2) \\ \lambda_1 & \lambda_2 \\ \theta_1 & \theta_2 \end{pmatrix} T_{0.6},$$

which implies  $\begin{pmatrix} h(p_1) & h(p_2) \\ \lambda_1 & \lambda_2 \\ \theta_1 & \theta_2 \end{pmatrix} \gg \begin{pmatrix} h(p_1^*) & h(p_2^*) \\ \lambda_1^* & \lambda_2^* \\ \theta_1^* & \theta_2^* \end{pmatrix}$ . Under this set up,

$$\bar{F}_{Y_{1:2}}(2.8) = 0.9999999923, \bar{F}_{Y_{1:2}^*}(2.8) = 0.9999999918, \bar{F}_{Y_{1:2}}(5) = 0.8918294672, \bar{F}_{Y_{1:2}^*}(5) = 0.9248659543, \text{ which readily shows that } Y_{1:2}^* \not\leq_{st} Y_{1:2}.$$

The following theorem extends Theorem 3.12 when two sets of  $n$ -independent observations are from ELS distribution. The generalization is the direct of the Theorem 3.12 and Lemma 7 of Balakrishnan et al. (2018). So, the proof is omitted.

**Theorem 3.14.** *Let Assumption 3.1 hold. Suppose  $h : [0, 1] \rightarrow R_+$  is differentiable and strictly increasing concave function. Further, let  $\tilde{r}(u)$  and  $u\tilde{r}(u)$  are increasing in  $u$ . Then, for  $i = 1, \dots, n$  and T-transform matrix  $T_w$ , if  $\alpha_i = \alpha_i^* = \alpha$ , and  $(h(p), \lambda, \theta) \in S_n$ , we have that*

$$\begin{pmatrix} h(p_1^*) & h(p_2^*) & \cdots & h(p_n^*) \\ \lambda_1^* & \lambda_2^* & \cdots & \lambda_n^* \\ \theta_1^* & \theta_2^* & \cdots & \theta_n^* \end{pmatrix} = \begin{pmatrix} h(p_1) & h(p_2) & \cdots & h(p_n) \\ \lambda_1 & \lambda_2 & \cdots & \lambda_n \\ \theta_1 & \theta_2 & \cdots & \theta_n \end{pmatrix} T_w,$$

implies  $Y_{1:n}^* \geq_{st} Y_{1:n}$ .

**Theorem 3.15.** *Let Assumption 3.1 hold and  $T_{w_1}, \dots, T_{w_k}$  be T-transform matrices with same structures. Suppose,  $h : [0, 1] \rightarrow R_+$  is differentiable and strictly increasing concave function. Further, let  $\tilde{r}(u)$ , and  $u\tilde{r}(u)$  are strictly increasing in  $u$ . Then, if  $\alpha_i = \alpha_i^* = \alpha$ , and  $(h(\underline{p}), \underline{\lambda}, \underline{\theta}) \in S_n$ , we have that*

$$\begin{pmatrix} h(p_1^*) & h(p_2^*) & \cdots & h(p_n^*) \\ \lambda_1^* & \lambda_2^* & \cdots & \lambda_n^* \\ \theta_1^* & \theta_2^* & \cdots & \theta_n^* \end{pmatrix} = \begin{pmatrix} h(p_1) & h(p_2) & \cdots & h(p_n) \\ \lambda_1 & \lambda_2 & \cdots & \lambda_n \\ \theta_1 & \theta_2 & \cdots & \theta_n \end{pmatrix} T_{w_1} \cdots T_{w_k},$$

implies  $Y_{1:n}^* \geq_{st} Y_{1:n}$ .

**Proof:** Since, a finite product of T-transform matrices with the same structure is also a T-transform matrix, so, the desired result can be obtained by repeating the result of Theorem 3.13.  $\square$

**Theorem 3.16.** *Let Assumption 3.1 hold and  $T_{w_1}, \dots, T_{w_k}$ ,  $k > 2$  be T-transform matrices with different structures. Suppose,  $h : [0, 1] \rightarrow R_+$  is differentiable and strictly increasing concave function. Further, let  $\tilde{r}(u)$  and  $u\tilde{r}(u)$  are increasing in  $u$ . Then, if  $\alpha_i = \alpha_i^* = \alpha$ , and  $(h(\underline{p}), \underline{\lambda}, \underline{\theta}) \in S_n$  and  $(h(\underline{p}), \underline{\lambda}, \underline{\theta})T_{w_1} \cdots T_{w_i} \in S_n, i = 1, \dots, k - 1$ , we have*

$$\begin{pmatrix} h(p_1^*) & h(p_2^*) & \cdots & h(p_n^*) \\ \lambda_1^* & \lambda_2^* & \cdots & \lambda_n^* \\ \theta_1^* & \theta_2^* & \cdots & \theta_n^* \end{pmatrix} = \begin{pmatrix} h(p_1) & h(p_2) & \cdots & h(p_n) \\ \lambda_1 & \lambda_2 & \cdots & \lambda_n \\ \theta_1 & \theta_2 & \cdots & \theta_n \end{pmatrix} T_{w_1} \cdots T_{w_k},$$

implies  $Y_{1:n}^* \geq_{st} Y_{1:n}$ .

**Proof:**

$$\begin{pmatrix} h^{(i)}(p_1) & \cdots & h^{(i)}(p_n) \\ \lambda_1^{(i)} & \cdots & \lambda_n^{(i)} \\ \theta_1^{(i)} & \cdots & \theta_n^{(i)} \end{pmatrix} = \begin{pmatrix} h(p_1) & \cdots & h(p_n) \\ \lambda_1 & \cdots & \lambda_n \\ \theta_1 & \cdots & \theta_n \end{pmatrix} T_{w_1} \cdots T_{w_i}, \text{ for } i = 1, \dots, k.$$

Assume  $V_1^{(i)}, \dots, V_n^{(i)}$ ,  $i = 1, \dots, k$ , are independent sets of random variables with  $V_j^{(i)} \sim ELS(\lambda_j^{(i)}, \theta_j^{(i)}, \alpha_j)$  where  $\alpha_i = \alpha_i^* = \alpha$ ,  $j = 1, \dots, n$  and  $i = 1, \dots, k$ . From the assumption of the theorem, it follows that

$$\begin{pmatrix} h(p_1)^{(i)} & \cdots & h(p_n)^{(i)} \\ \lambda_1^{(i)} & \cdots & \lambda_n^{(i)} \\ \theta_1^{(i)} & \cdots & \theta_n^{(i)} \end{pmatrix} \in S_n.$$

using the results of Theorem 3.13, it then follows that  $Y_{1:n} \leq_{st} V_{1:n}^{(1)} \leq_{st} \cdots \leq_{st} V_{1:n}^{(k-2)} \leq_{st} V_{1:n}^{(k-1)} \leq_{st} Y_{1:n}^*$ . Which completes the proof of the theorem.  $\square$

**Theorem 3.17.** *Let Assumption 3.1 hold for  $n = 2$ . Suppose  $h : [0, 1] \rightarrow R_+$  is a differentiable and strictly increasing concave function. Further, assume that  $\tilde{r}(u)$  and  $u\tilde{r}(u)$  are increasing in  $u$ , and  $(h(p), \lambda, \theta, \alpha) \in U_2$ . Then,*

$$\begin{pmatrix} h(p_1) & h(p_2) \\ \lambda_1 & \lambda_2 \\ \theta_1 & \theta_2 \\ \alpha_1 & \alpha_2 \end{pmatrix} \gg \begin{pmatrix} h(p_1^*) & h(p_2^*) \\ \lambda_1^* & \lambda_2^* \\ \theta_1^* & \theta_2^* \\ \alpha_1^* & \alpha_2^* \end{pmatrix},$$

implies  $Y_{1:2}^* \geq_{st} Y_{1:2}$ .

**Proof:** With the help of Lemma 2.1, the proof follows from arguments similar to those in the proof of Theorem 3.13. It is omitted for brevity.  $\square$

We present a counterexample to show that the comparison result may not hold if the assumptions are not satisfied.

**Counterexample 3.3** Let the baseline distribution function be  $F(t) = 1 - \exp(1 - t^{0.5})$ ,  $t \geq 1$ . Here  $\tilde{r}(t)$  and  $t\tilde{r}(t)$  are decreasing. Take  $h(p) = e^p$ , where  $h(p)$  is convex. Thus, the assumption of Theorem 3.17 are not violated. Let us set  $(\alpha_1, \alpha_2) = (0.2, 0.5)$ ,  $(\alpha_1^*, \alpha_2^*) = (0.44, 0.26)$ ,  $(\lambda_1, \lambda_2) = (1, 1.5)$ ,  $(\lambda_1^*, \lambda_2^*) = (1.4, 1.1)$ ,  $(\theta_1, \theta_2) = (4, 2)$ ,  $(\theta_1^*, \theta_2^*) = (2.4, 3.6)$ ,  $(p_1, p_2) = (\ln(4), \ln(5))$ ,  $(p_1^*, p_2^*) = (\ln(4.8), \ln(4.2))$ . Consider a T-transform matrix  $T_{0.2} = \begin{pmatrix} 0.2 & 0.8 \\ 0.8 & 0.2 \end{pmatrix}$ . Then, it can be shown that

$$\begin{pmatrix} h(p_1^*) & h(p_2^*) \\ \lambda_1^* & \lambda_2^* \\ \theta_1^* & \theta_2^* \\ \alpha_1^* & \alpha_2^* \end{pmatrix} = \begin{pmatrix} h(p_1) & h(p_2) \\ \lambda_1 & \lambda_2 \\ \theta_1 & \theta_2 \\ \alpha_1 & \alpha_2 \end{pmatrix} T_{0.2},$$

Which implies  $\begin{pmatrix} h(p_1) & h(p_2) \\ \lambda_1 & \lambda_2 \\ \theta_1 & \theta_2 \\ \alpha_1 & \alpha_2 \end{pmatrix} \gg \begin{pmatrix} h(p_1^*) & h(p_2^*) \\ \lambda_1^* & \lambda_2^* \\ \theta_1^* & \theta_2^* \\ \alpha_1^* & \alpha_2^* \end{pmatrix}$ . Finally  $\bar{F}_{Y_{1:2}}(5) -$

$\bar{F}_{Y_{1:2}^*}(5) = 0.4133022299$ ,  $\bar{F}_{Y_{1:2}}(7.5) - \bar{F}_{Y_{1:2}^*}(7.5) = -0.0324688838$ , which readily shows that  $Y_{1:2}^* \not\geq_{st} Y_{1:2}$ .

In the following theorem, we present a generalization of Theorem 3.17 to the case of  $n$  independent variables.

**Theorem 3.18.** *Let Assumption 3.1 hold. Further, let  $T_w$  be a T-transform matrix. Suppose  $h : [0, 1] \rightarrow R_+$  is a differentiable strictly increasing concave function. Further, assume that  $\tilde{r}(u)$  and  $u\tilde{r}(u)$  are increasing in  $u$ , and*



let  $(h(\underline{p}), \underline{\lambda}, \underline{\theta}, \underline{\alpha}) \in U_n$ . Then,

$$\begin{pmatrix} h(p_1^*) & h(p_2^*) & \cdots & h(p_n^*) \\ \lambda_1^* & \lambda_2^* & \cdots & \lambda_n^* \\ \theta_1^* & \theta_2^* & \cdots & \theta_n^* \\ \alpha_1^* & \alpha_2^* & \cdots & \alpha_n^* \end{pmatrix} = \begin{pmatrix} h(p_1) & h(p_2) & \cdots & h(p_n) \\ \lambda_1 & \lambda_2 & \cdots & \lambda_n \\ \theta_1 & \theta_2 & \cdots & \theta_n \\ \alpha_1 & \alpha_2 & \cdots & \alpha_n \end{pmatrix} T_w,$$

implies  $Y_{1:n}^* \geq_{st} Y_{1:n}$ .

**Theorem 3.19.** Let  $T_{w_1}, \dots, T_{w_k}$  be T-transform matrices with same structure. Let Assumption 3.1 hold and  $h : [0, 1] \rightarrow R_+$  be a differentiable and strictly increasing concave function. Further, assume that  $\tilde{r}(u)$  and  $u\tilde{r}(u)$  are strictly increasing in  $u$ , and  $(h(\underline{p}), \underline{\lambda}, \underline{\theta}, \underline{\alpha}) \in U_n$ . Then,

$$\begin{pmatrix} h(p_1^*) & h(p_2^*) & \cdots & h(p_n^*) \\ \lambda_1^* & \lambda_2^* & \cdots & \lambda_n^* \\ \theta_1^* & \theta_2^* & \cdots & \theta_n^* \\ \alpha_1^* & \alpha_2^* & \cdots & \alpha_n^* \end{pmatrix} = \begin{pmatrix} h(p_1) & h(p_2) & \cdots & h(p_n) \\ \lambda_1 & \lambda_2 & \cdots & \lambda_n \\ \theta_1 & \theta_2 & \cdots & \theta_n \\ \alpha_1 & \alpha_2 & \cdots & \alpha_n \end{pmatrix} T_{w_1} \cdots T_{w_k},$$

implies  $Y_{1:n}^* \geq_{st} Y_{1:n}$ .

The following theorem presents a generalization to the case of a finite number of T-transform matrices with different structures.

**Theorem 3.20.** Let Assumption 3.1 hold. Further, let  $T_{w_1}, \dots, T_{w_k}$ ,  $k > 2$  be T-transform matrices, with different structures. Suppose  $h : [0, 1] \rightarrow R_+$  is a differentiable and strictly increasing concave function. Further, assume that  $\tilde{r}(u)$  and  $u\tilde{r}(u)$  are increasing in  $u$ , and  $(h(\underline{p}), \underline{\lambda}, \underline{\theta}, \underline{\alpha}) \in U_n$  and  $(h(\underline{p}), \underline{\lambda}, \underline{\theta}, \underline{\alpha})T_{w_1} \cdots T_{w_i} \in U_n, i = 1, \dots, k - 1$ . Then,

$$\begin{pmatrix} h(p_1^*) & h(p_2^*) & \cdots & h(p_n^*) \\ \lambda_1^* & \lambda_2^* & \cdots & \lambda_n^* \\ \theta_1^* & \theta_2^* & \cdots & \theta_n^* \\ \alpha_1^* & \alpha_2^* & \cdots & \alpha_n^* \end{pmatrix} = \begin{pmatrix} h(p_1) & h(p_2) & \cdots & h(p_n) \\ \lambda_1 & \lambda_2 & \cdots & \lambda_n \\ \theta_1 & \theta_2 & \cdots & \theta_n \\ \alpha_1 & \alpha_2 & \cdots & \alpha_n \end{pmatrix} T_{w_1} \cdots T_{w_k},$$

implies  $Y_{1:n}^* \geq_{st} Y_{1:n}$ .

---

#### 4. SOME SPECIAL CASES

---

In this section, we present some special cases of the results obtained in the previous sections. We consider two special distributions - generalized gamma, half - normal distributions. For these distributions, we present some comparisons results using the general results established earlier. In terms of hazard rate function, the following theorems can be proved like the theorems above, which are proved by reversed hazard rate function. To prevent recurrence, you can refer to Das et al. (2021).

---

#### 4.1. Generalized gamma distribution

---

In this subsection, we consider the generalized gamma distribution with density function

$$f(t) \propto t^{a-1} e^{-t^b}, a, b, t > 0.$$

It is easy to check that the hazard rate function of this distribution is increasing for  $a, b \geq 1$  and decreasing for  $0 < a, b \leq 1$  (see Hazra et al., 2018). We now consider the baseline distribution function to the generalized gamma distribution.

**Theorem 4.1.** For a baseline distribution function  $F(\cdot)$ , let  $X_1, \dots, X_n$  ( $X_1^*, \dots, X_n^*$ ) be non-negative independent random variables with  $X_i \sim ELS(\lambda_i, \theta_i, \alpha_i)$  [ $X_i^* \sim ELS(\mu_i, \delta_i, \beta_i)$ ],  $i = 1, \dots, n$ . Further, let  $I_{p_1}, \dots, I_{p_n}$  [ $I_{p_1}^*, \dots, I_{p_n}^*$ ] be a set of independent Bernoulli random variables, independent of  $X_i$  [ $X_i^*$ ]'s with  $E(I_{p_i}) = p_i$  [ $E(I_{p_i}^*) = p_i^*$ ],  $i = 1, \dots, n$ . Further, let  $h : [0, 1] \rightarrow R_+$  be a differentiable, increasing and convex function. Then, for  $i = 1, 2$ , if  $\theta_i = \delta_i = \delta$  and  $\alpha_i = \beta_i = \alpha \geq 1$ ,  $r(x)$  is increasing. Suppose  $h(p) = e^p \ln(1 + p)$ . Then, for  $a, b \geq 1$ , we have  $\begin{pmatrix} h(p_1) & h(p_2) \\ \theta_1 & \theta_2 \end{pmatrix} \gg \begin{pmatrix} h(p_1^*) & h(p_2^*) \\ \delta_1 & \delta_2 \end{pmatrix}$  implies  $Y_{1:2}^* \geq_{st} Y_{1:2}$ .

**Theorem 4.2.** For a baseline distribution function  $F(\cdot)$ , let  $X_1, \dots, X_n$  ( $X_1^*, \dots, X_n^*$ ) be non-negative independent random variables with  $X_i \sim ELS(\lambda_i, \theta_i, \alpha_i)$  [ $X_i^* \sim ELS(\mu_i, \delta_i, \beta_i)$ ],  $i = 1, \dots, n$ . Further, let  $I_{p_1}, \dots, I_{p_n}$  [ $I_{p_1}^*, \dots, I_{p_n}^*$ ] be a set of independent Bernoulli random variables, independent of  $X_i$  [ $X_i^*$ ]'s with  $E(I_{p_i}) = p_i$  [ $E(I_{p_i}^*) = p_i^*$ ],  $i = 1, \dots, n$ . Further, let  $h : [0, 1] \rightarrow R_+$  be a differentiable, increasing and convex function. Further, let the baseline hazard rate  $r(\cdot)$  be increasing. If  $\alpha_i = \beta_i = \alpha \geq 1$  and  $(h(p), \lambda, \theta) \in N_2$ . Suppose  $h(p) = p^2$ .

Then, for  $a, b \geq 1$ , we have  $\begin{pmatrix} h(p_1) & h(p_2) \\ \lambda_1 & \lambda_2 \\ \theta_1 & \theta_2 \end{pmatrix} \gg \begin{pmatrix} h(p_1^*) & h(p_2^*) \\ \mu_1 & \mu_2 \\ \delta_1 & \delta_2 \end{pmatrix}$  implies  $Y_{1:2}^* \geq_{st} Y_{1:2}$ .

---

#### 4.2. Half-normal distribution

---

Consider the probability distribution function of a half-normal distribution given by

$$f(t) \propto e^{-\frac{t^2}{2}}, t > 0.$$

The hazard rate function of the above half-normal distribution is increasing (see Hazra et al. (2018)). The distribution function of the half-normal distribution is now taken as the baseline distribution function.

**Theorem 4.3.** For a baseline distribution function  $F(\cdot)$ , let  $X_1, \dots, X_n$  ( $X_1^*, \dots, X_n^*$ ) be non-negative independent random variables with  $X_i \sim ELS(\lambda_i, \theta_i, \alpha_i)$  ( $X_i^* \sim ELS(\mu_i, \delta_i, \beta_i)$ ),  $i = 1, \dots, n$ . Further, let  $I_{p_1}, \dots, I_{p_n}$  [ $I_{p_1}^*, \dots, I_{p_n}^*$ ] be a set of independent Bernoulli random variables, independent of  $X_i$  [ $X_i^*$ ]'s with  $E(I_{p_i}) = p_i$  [ $E(I_{p_i}^*) = p_i^*$ ],  $i = 1, \dots, n$ . Further, let  $h : [0, 1] \rightarrow R_+$  be a differentiable, increasing and convex function. Further, let the baseline hazard rate  $r(\cdot)$  be increasing. If  $\theta_i = \delta_i = \theta$  and  $(h(p), \lambda, \alpha) \in N_2^*$ . Let  $h(p) = -p \ln(1 -$

$p)$ . Then  $\begin{pmatrix} h(p_1) & h(p_2) \\ \lambda_1 & \lambda_2 \\ \alpha_1 & \alpha_2 \end{pmatrix} \gg \begin{pmatrix} h(p_1^*) & h(p_2^*) \\ \mu_1 & \mu_2 \\ \beta_1 & \beta_2 \end{pmatrix}$  implies  $Y_{1:2}^* \geq_{st} Y_{1:2}$ .

---

## 5. APPLICATIONS

---

In this section, we discuss application of few of our established results in insurance and auction theory. Suppose  $X_1, \dots, X_n$  are independent exponentiated location-scale random variables with  $X_i \sim ELS(\lambda_i, \mu_i, \alpha_i)$ , for  $i = 1, \dots, n$ , and  $I_{p_1}, \dots, I_{p_n}$  are independent Bernoulli random variables, independent of the  $X_i$ 's, with  $E(I_{p_i}) = p_i$ . Let  $Y_i = I_{p_i}X_i$ , for  $i = 1, \dots, n$ . Suppose  $X_i$  denotes the total of random claims that can be made in an insurance period and  $I_{p_i}$  denotes a Bernoulli random variables associated with  $X_i$  defined as follows:  $I_{p_i} = 1$  when ever the  $i$ th policyholder makes random claim  $X_i$  and  $I_{p_i} = 0$  whenever he/she does not make a claim. In setting,  $Y_i = I_{p_i}X_i$  corresponds to the claim amount in a portfolio of risks. The problem of comparison of number of claims and aggregate claim amounts with respect to some well-known stochastic orders is of interest on both theoretical and practical view points. Under some conditions Theorems 3.2, 3.3, 3.4 respectively conclude that  $Y_{n:n}$  in the weakly supermajorized order, weakly submajorized order,  $p$ -larger and reciprocally majorized order is stochastically smaller. There are many real-life applications of the ordering results. We discuss applications of few of our established results in auction theory. Auction theory has been an interest topic to various scientists because of its usefulness for sale of variety of items or purchasing services. For more details in auction theory, we refer to (Klemperer, (2004)). In real world, among all types of auctions, the sealed-bid private-value auction is of theoretical interest. Also, this type of auction has been used extensively. In this case, bidders hand in their bids to the auctioneer simultaneously and can neither observe their rival bids nor revise their own bids. The bidders having the highest bid wins. The bidders with the lowest bid wins in the reverse auction. Consequently, the bidder pays his own bid in the sealed-bid first-price auction (FPA). Few of our established results could be useful for some new light in the auction theory. Let the bids follow exponentiated locatio-scale model. Then, under some conditions, Theorems 3.2, 3.3, 3.4 respectively conclude that the final price in the FPA with more heterogeneous shape parameters in the weakly super majorized order, reciprocal of the shape parameters in the weakly submajorized, scale parameters in the  $p$ -larger and reciprocally majorized orders is stochastically smaller.

---

## 6. CONCLUDING

---

In this paper, when the matrix of parameters changes to another matrix of parameters with respect to multivariate chain majorization, we study the usual stochastic order of the smallest order statistics when each component receives a random shock. Under certain conditions, by using the concept of vector majorization and related orders, we have also discussed stochastic comparison between series and parallel systems in the sense of the usual stochastic order under random shock. We have then applied the results for some special cases of the exponentiated location-scale model with possibly different scale, location and shape parameters to illustrate the established results.

---

## ACKNOWLEDGMENTS

---

The authors are thankful to the Editor and an anonymous referee for some useful comments and suggestions, which have resulted in improving the content and the presentation of the paper.

---

## REFERENCES

---

- [1] BALAKRISHNAN, N.; HAIDARI, A. and MASOUMIFARD, K. (2015). Stochastic comparisons of series and parallel systems with generalized exponential components, *IEEE Transaction on Reliability*, **64**, 1, 333-348.
- [2] BALAKRISHNAN, N.; NANDA, P. and KAYAL, S. (2018). Ordering of series and parallel systems comprising heterogeneous generalized modified Weibull components, *Applied Stochastic Models in Business and Industry*, **34**, 6, 816-834.
- [3] BARMALZAN, G.; NAJAFABADI, A.T.P. and BALAKRISHNAN, N (2017). Ordering properties of the smallest and largest claim amounts in a general scale model, *Scandinavian Actuarial Journal* **2017**, 2, 105-124.
- [4] CHOWDHURY, S. and KUNDU, A. (2017). Stochastic comparisons of parallel systems with log-Lindley distributed components, *Operations Research Letters* **45**, 3, 199-205.
- [5] CHOWDHURY, S. and KUNDU, A. (2018). Stochastic comparisons of parallel systems with log-Lindley distributed components under random shocks, arXiv preprint arXiv: 1804.04103.
- [6] CHOWDHURY, S.; KUNDU, A. and MISHRA, S. K. (2021). On comparison of two parallel systems having Log-Lindley distributed components, *ommunication in Statistics - Theory and Methods* <http://doi.org/10.1080/03610926.2021.1910838>.

- [7] DAS, S.; KAYAL, S. and CHOWDHURY, D. (2021). Ordering results on extremes of exponentiated location-scale models, *Probability in the Engineering and Informational Sciences*, **35**, 2, 331-354.
- [8] DAS, S. and KAYAL, S. (2021). Ordering extremes of exponentiated location-scale models with dependent and heterogeneous random samples, *Metrika*, <https://doi.org/10.1007/s00184-019-00753-2>.
- [9] DYKSTRA, R.; KOCHAR, SC. and ROJO, J. (1997). Stochastic comparisons of parallel systems of heterogeneous exponential components, *Journal of Statistical Planning and Inference*, **65**, 203-211.
- [10] FANG, L. and BALAKRISHNAN, N. (2018). Ordering properties of the smallest order statistics from generalized Birnbaum-Saunders models with associated random shocks, *Metrika*, **81**, 1, 19-35.
- [11] FANG, R.; LI, C. and LI, X. (2016). Stochastic comparisons on sample extremes of dependent and heterogeneous observations, *Statistics*, **50**, 4, 930-955.
- [12] FANG, L. and ZHANG, X. (2013). Stochastic comparisons of series systems with heterogeneous Weibull components, *Statistic and Probability Letters*, **83**, 7, 1649-1653.
- [13] FANG, L. and XU, T. (2019). Ordering results of the smallest and largest order statistics from independent heterogeneous exponentiated gamma random variables, *Statistica Neerlandica*, **73**, 2, 197-210.
- [14] HAZRA, N.K.; KUITI, M.R.; FINKELSTEIN, M. and NANDA, A.K. (2017). On stochastic comparisons of maxima order statistics from the location-scale family of distributions, *Journal of multivariate Analysis*, **16**, 31-41.
- [15] HAZRA, N.K.; KUITI, M.R.; FINKELSTEIN, M. and NANDA, A.K. (2018). On stochastic comparisons of minimum order statistics from the location-scale family of distributions, *Metrika*, **81**, 2, 105-123.
- [16] KHALEDI, B.E. and KOCHAR, S.C. (2002). Dispersive ordering among linear combination of uniform random variables, *Journal of Statistical planning and Inference*, **100**, 1, 13-21.
- [17] KLEMPERER, P. (2004). *Auction: Theory and Practice*, New York: Academic Press.
- [18] KOCHAR, S.C. and TORRADO, N. (2015). On stochastic comparisons of largest order statistics in the scale model, *Communications in Statistics-Theory and Methods*, **44**, 19, 4132-4143.
- [19] KUNDU, A. and CHOWDHURY, S. (2016). Ordering properties of order statistics from heterogeneous exponentiated Weibull models, *Statistics and Probability Letters*, **114**, 119-127.
- [20] KUNDU, A. and CHOWDHURY, S. (2018). Ordering properties of sample minimum from Kumaraswamy-G random variables, *Statistics*, **52**, 1, 133-146.
- [21] KUNDU, A. and CHOWDHURY, S. (2019). Ordering properties of the largest order statistics from Kumaraswamy-G models under random shocks, *Communication in Statistics - Theory and Method*, **50**, 6, 1502-1514.
- [22] KUNDU, A. and CHOWDHURY, S. (2020). On stochastic comparisons of series systems with heterogeneous dependent and independent location-scale family distributed components, *Operation Research Letters*, **48**, 1, 40-47.

- [23] KUNDU, A.; CHOWDHURY, S. and BALAKRISHNAN, N. (2021). Ordering properties of the smallest and largest lifetimes in Gompertz-Makeham model, *Communication in Statistics - Theory and Methods*, DOI: 10.1080/03610926.2021.1919898.
- [24] KUNDU, A.; CHOWDHURY, S.; NANDA, A. K. and HAZRA, N. K. (2016). Some results on majorization and their applications, *Journal of Computational and Applied Mathematics*, **301**, 161-177.
- [25] LI, C. and LI, X. (2015). Likelihood ratio order of sample minimum from heterogeneous Weibull random variables, *Statistics and Probability Letters*, **97**, 46-53.
- [26] MARSHALL A.W. and OLKIN I. (1997). A new method for adding a parameter to a family of distributions with application to the exponential and Weibull families, *Biometrika*, **84**, 641-652.
- [27] MARSHALL, A.W.; OLKIN, I. and ARNOLD, B.C. (2011). *Inequalities: Theory of Majorization and Its Applications*, Springer series in Statistics, New York.
- [28] TORRADO, N. (2015). On magnitude ordering between smallest order statistics from heterogeneous beta distributions, *Journal of Mathematical Analysis and Applications*, 824-838.
- [29] TORRADO, N. and KOCHAR, S.C. (2015). Stochastic order relations among parallel systems from Weibull distributions, *Journal of Applied Probability*, **52**, 1, 102- 116.
- [30] ZHAO, P. and BALAKRISHNAN, N. (2011). Some characterization results for parallel systems with two heterogeneous exponential components, *Statistics*, **45**, 6, 593-604.